arXiv:1507.07329v2 [math.AP] 31 Jul 2015

A boundary partial regularity and a regularity criterion for New Harmonic Heat flows

Kazuhiro HORIHATA

August 3, 2015

Abstract

In my previous paper of K.Horihata [9], we have proposed a Ginzburg-Landau type heat flow with a time-dependent parameter and then passing to the limit of the parameter appeared in it, we have constructed a harmonic heat flow into spheres. By using this scheme, we establish a few energy inequalities:

(i) monotonical inequalities and

(ii) a reverse Poincare inequality at any boundary point.

Furthermore we assert that these inequalities (i) and (ii) derive the smaller estimates on the set of noncontinuous points for our flow contrast to the former results. We refer to them by Y.Chen [2] and Y.Chen, J.Li, F.H.Lin [4]. Next we introduce two conditions for the whole domain's regularity for it; The one is a boundary energy smallness and the another is an one-sided condition proposed by S.Hildebrandt and K.-O.Widman [8].

1 Introduction.

A previous paper: K.Horihata [9], constructed a harmonic heat flow between d-dimensional unit ball and D-dimensional unit sphere with

(i) a global energy inequality,

(ii) a monotonical inequality

(iii) a reverse Poincaré inequality

and showed the flow is smooth except a small singular set. This paper discusses a partial regularity for such a flow from C^2-domain in $\mathbb{R}^d$ to the D-dimensional unit sphere and then state two condition for the whole domain's smoothness for it under a certain hypothesis: For any point $x_0 \in \partial\Omega$, there exist a neighbourhood $U(x_0)$ and C^2-diffeomorphism Φ_{x_0} between $U(x_0)$ and $\mathbb{R}^d$, writing

$$\begin{gathered} y \;=\; (y', y_d) \;=\; \Phi_{x_0}(x) \;=\; (x', x_d \,-\, \phi_{x_0}(x')) \\ (y \;=\; (y', y_d) \;\in\; \mathbb{R}^d,\; x \;=\; (x', x_d) \;\in\; U(x_0)) \end{gathered}$$

whereupon writing $x_0 = (x'_0, x_{d,0})$, a suitable C^2-function ϕ_{x_0} on $U(x_0) \cap \mathbb{R}^d_0$ satisfies

$$\begin{aligned} &\phi_{x_0}(x'_0) \;=\; 0,\; \nabla\phi_{x_0}(x'_0) \;=\; 0, \\ &\sum_{i,j=1}^{d-1} \nabla_i\nabla_j\phi_{x_0}(x'_0)(x' - x'_0)_i(x' - x'_0)_j \;\geq\; \theta_0|x' - x'_0|^2 \end{aligned} \tag{B}$$

for any point $x = (x', x_d) \in U(x_0) \cap \partial\Omega$, where a positive constant θ_0 is independent of x_0. We call the condition our the boundary (B).

We rigorously state our problem and results: Let Ω and $\mathbb{S}^D$ be respectively the C^2-domain in $\mathbb{R}^d$ (*not always bounded*) and the unit sphere in $\mathbb{R}^{D+1}$ and a parabolic cylinder $Q(T)$ be $(0, T) \times \Omega$, where d and D are positive integers greater than or equal to 2. Consider the Sobolev class $H^{k,p}_B(\Omega; \mathbb{S}^D)$ $:= \{u \in L(\Omega \cap B_n(0); \mathbb{R}^{D+1}),\ \nabla u \in H^{k-1,p}\ (\Omega; \mathbb{R}^{D+1})\ ; |u| = 1 \quad \text{a.e. } x \in \Omega\}$. for any positive integer n, where k is any nonnegative and p is positive integers greater than equal to 1. Giving a mapping $u_0 \in H^{1,2}_B(\Omega; \mathbb{S}^D)$, we consider the heat flow:

$$\left\{ \begin{array}{rcll} \dfrac{\partial u}{\partial t} & = & \triangle u \,+\, |\nabla u|^2 u & \text{in } \ Q(T), \\ u(0, x) & = & u_0(x) & \text{at } \quad \{0\} \times \Omega, \\ u(t, x) & = & u_0(x) & \text{on } \quad [0, T) \times \partial\Omega. \end{array} \right. \tag{1.1}$$

Define three function classes:

$$\begin{aligned} & L^\infty\big(0, T; H^{1,2}_B(\Omega; \mathbb{S}^D)\big) \\ & \qquad := \{u | u \text{ is measurable from } [0, T] \text{ to } H^{1,2}(\Omega; \mathbb{S}^D) \\ & H^{1,2}\big(0, T; L^2_B(\Omega; \mathbb{R}^{D+1})\big) \\ & \qquad := \Big\{u \;\in\; L^2_B(\Omega; \mathbb{R}^{D+1})\,;\, \partial u/\partial t \;\in\; L^2(Q(T); \mathbb{R}^{D+1})\Big\} \\ & \quad V_B(Q(T); \mathbb{S}^D) \;:=\; L^\infty\big(0, T; H^{1,2}_B(\Omega; \mathbb{S}^D)\big) \cap H^{1,2}\big(0, T; L^2_B(\Omega; \mathbb{R}^{D+1})\big). \end{aligned}$$

The weak formulation of (1.1) is as follows: For any given mapping $u_0 \in H^{1,2}_B(\Omega; \mathbb{S}^D)$, we call a mapping $u \in V_B(Q(T); \mathbb{S}^D)$ a weakly harmonic heat flow (*WHHF*) provided for any $\phi \in C_0^\infty(Q(T); \mathbb{R}^{D+1})$

$$\int\limits_{Q(T)} \Big(\langle \frac{\partial u}{\partial t}, \phi \rangle \, + \, \langle \nabla u, \nabla \phi \rangle - \langle u, \phi \rangle |\nabla u|^2 \Big) \, dz \, = \, 0, \tag{1.2}$$

$$u(t) \, - \, u_0 \in \overset{\circ}{H}{}^{1,2}_B(\Omega; \mathbb{R}^{D+1}) \qquad \text{for almost every } t \, \in \, (0,T), \tag{1.3}$$

$$\lim_{t \searrow 0} u(t) \, = \, u_0 \qquad\qquad \text{in} \quad L^2_B(\Omega; \mathbb{R}^{D+1}). \tag{1.4}$$

K.Horihata [9] has proposed a new approximate evolutional scheme said to be the Ginzburg-Landau heat flow (*GLHF*). We review Ginzburg-Landau heat flow: Introduce smooth functions $\chi(t)$ and $\kappa(t)$ by

$$\begin{aligned} \chi(t) \, &= \, \begin{cases} t & (t \, < \, 2) \\ 3 & (t \, \geq \, 4), \end{cases} \\ \kappa(t) \, &= \, \arctan(t)/\pi. \end{aligned} \tag{1.5}$$

Let a mapping u_0 in $H^{1,2}_B \, (\Omega \, ; \, \mathbb{S}^D) \cap (H_B^{2-1/q,q} \cap H_B^{[(d+1)/2]+1,p}) \, (\Omega \setminus (\complement\Omega)_{\delta_0}; \mathbb{S}^D)$ with positive numbers sufficiently small δ_0, and positive numbers q greater than or equal to $d+2$ and p greater than 2. Then GLHF is designated by solution of the systems:

$$\frac{\partial u_\lambda}{\partial t} \, - \, \triangle u_\lambda \, + \, \lambda^{1-\kappa} \dot{\chi}\big((|u_\lambda|^2 \, - \, 1)^2 \big) (|u_\lambda|^2 \, - \, 1) u_\lambda \, = \, 0 \quad \text{in} \quad Q(T), \tag{1.6}$$

$$u_\lambda \, = \, u_0 \quad \text{on} \quad \partial Q(T). \tag{1.7}$$

If you notice that the nonlinear term of $\lambda^{1-\kappa} \, \dot{\chi}\big((|u_\lambda|^2 \, - \, 1)^2 \big) \, \big(|u_\lambda|^2 \, - \, 1 \big) u_\lambda$ is bounded, Banach's fixed point theorem can state the unique existence of the mapping u_λ on $Q(T)$ with

(a) $u_\lambda \, \in \, C^\infty(Q(T))$,

(b) (1.8) is fulfilled in $Q(T)$,

(c) $u_\lambda(t,x) \, - \, u_0(x) \in \overset{\circ}{H}{}^{1,2}_B(\Omega; \mathbb{R}^{D+1})$ for almost every t in $(0,T)$,

(d) $\lim_{t \searrow 0} ||u_\lambda(t, \cdot) \, - \, u_0(\cdot)||_{L^2_B(\Omega)} \, = \, 0$.

An observation by K.Horihata [9, Theorem 2.4] and an application of a maximal principle on an unbounded domain which referred to F.John [10, Chap 7] prevails $|u_\lambda| \leq 1$ in $Q(T)$. Thus we rewrite (1.6) to

$$\frac{\partial u_\lambda}{\partial t} - \triangle u_\lambda + \lambda^{1-\kappa}\big(|u_\lambda|^2 - 1\big)u_\lambda = 0 \quad \text{in} \quad Q(T). \tag{1.8}$$

The benefits of my scheme is to easily derive

$$\int_{Q(T)} \lambda^{1-\kappa}(|u_\lambda|^2 - 1)^2 \, dz = O(1/\log\lambda) \quad \text{as } \lambda \nearrow \infty.$$

We outlook the history on a boundary regularity of harmonic mapping or harmonic heat flows: For the energy minimizing map, R.Schoen and K.Uhlenbeck [16] has established a boundary regularity for an appropriate boundary condition. A similar results on minimizers of a certain functional can be found in J.Jost and M.Meier [11]. J.Qing [14] proved the boundary regularity of weakly harmonic maps from surfaces with the boundary. C.Poon [13] constructed a smooth harmonic map u_a between B^3 and $\mathbb{S}^2$ except a prescribed point $a \in \overline{B}^3$, noncontinuous at a and $u_a = x$ on ∂B^3.

On the other hand, the papers discussed boundary regularity results on a harmonic heat flow are not so many. We refer it to Y.Chen [2] or Y.Chen-F.-H.Lin [3] or Y.Chen-J.Li-F.H.Lin [4] or C.Y.Wang [17]. Since the maps u_a by C.Poon [13] is also a non-smooth weakly harmonic heat flow, it may be reasonable to discuss a partial regularity for our harmonic heat flow and a regularity under a certain imposition.

This paper has two folds: The first constructs a few energy inequalities on $\overline{\Omega}$ and next discusses a partial regularity for our harmonic heat flow near the boundary; The proof proceeds as in the one of the author's former paper of K.Horihata [9] combined with the one by Y.Chen [2]. Thereafter by utilizing the reverse Poincaré inequality, we will prove that the WHHF is actually smooth except on a small set called "**singular set.**" More precisely we assert

Theorem 1.1 (Partial Regularity Theorem). *Let d be a positive integer larger than 2. For a mapping $u_0 \in H_B^{1,2}\,(\Omega\,;\mathbb{S}^D) \cap (H_B^{2-1/q,q} \cap H_B^{[(d+1)/2]+1,p})$ $(\Omega \setminus (\complement\Omega)_{\delta_0};\mathbb{S}^D)$ with a positive number sufficiently small δ_0, and positive numbers q and p respectively greater than or equal to $d+2$ and greater than 2, there exists a WHHF and it is smooth on a certain relative open set in $\overline{Q}(T)$ off a set called* **sing**. *The set has the finite $(d-\gamma_0)$-dimensional Hausdorff measure with respect to the parabolic metric, where γ_0 is a small positive* *number depending only on* u_0, d *and* $Q(T)$. *The WHHF also holds*

$$\int_{t_0-(2R_1)^2}^{t_0-R_1^2} \int_{\Omega} |\nabla u|^2 G_{z_0}\,dz + 2\int_{R_1}^{R_2} dR \int_{t_0-(2R)^2}^{t_0-R^2} dt \int_{\Omega} \Big|\frac{\partial u_\lambda}{\partial t} - \frac{x-x_0}{2\sqrt{t_0-t}}\cdot \nabla u\Big|^2 G_{z_0}\,dx$$

$$\leq C(R_2^{\mu_0} - R_1^{\mu_0}) \int_{t_0-(2R_2)^2}^{t_0-R_2^2} dt \int_{\Omega} |\nabla u|^2 G_{z_0}\,dx + C(R_2 - R_1) \tag{1.9}$$

for any positive numbers R_1 *and* R_2 *with* $R_1 \leq R_2$ *and an arbitrary point* $z_0 = (t_0, x_0)$ *in* $\overline{Q}(T)$ *satisfying* $R_2 < \sqrt{t_0/4}$, *where two positive constants* μ_0 *and* $C(\mu_0)$ *has the relation of* $C(\mu_0) \nearrow \infty$ *as* $\mu_0 \searrow 0$ *and in addition*

$$\int_{P_R(z_0)} |\nabla u(z)|^2\,dz \leq \frac{C}{R^2}\int_{P_{2R}(z_0)} |u(z) - h_0(x)|^2\,dz$$

$$+ \int_{P_R(z_0)} (|\nabla^{[(d+1)/2]+1} h_0(x)|^2 + |\nabla h_0(x)|^2)\,dz \tag{1.10}$$

for any parabolic cylinder $P_{2R}\,(z_0)$, *where a function* G_{z_0} *means a backward heat kernel indicated by*

$$G_{z_0}(t,x) = \frac{1}{\sqrt{4\pi(t_0-t)}^{\,d}} \exp\Big(-\frac{|x-x_0|^2}{4(t_0-t)}\Big) \quad ((t,x) \in (0,t_0)\times \mathbb{R}^d).$$

The subsequent theorems will take up two sufficient conditions on the whole domain's regularity for our *WHHF*. The one is the boundary energy smallness at a large time and the another is a target restriction on the initial mapping called "one-sided condition". We refer it to M.Giaquinta [6, p.237, Theorem 3.2].

Theorem 1.2 (First Regularity Theorem). *Assume that our domain is bounded with* (B) *and there exists some positive number* ϵ_0 *such that for all point* $z_0 = (t_0, x_0)$ *and some positive number* R_0 *less than* $\sqrt{t_0}/2$, *if we have*

$$\frac{\exp^{(4R_0)^{\mu_0}}}{R_0^2}\Big[\frac{\exp^{-4(d-2)/d_0^2}}{t_0^{(d-2)/2}} \int_{\Omega} |\nabla u_0|^2\,dx \tag{1.11}$$

$$+ \int_0^{t_0-R_0^2} dt \int_{\partial\Omega} |\nabla_\tau u_0|^2 G_{z_0}\big(d_{x_0} + \frac{4(t_0-t)}{d_0^2}\big)\,d\mathcal{H}_x^{d-1}\Big] + C(\mu_0)R_0 \leq \epsilon_0^2,$$

where μ_0 *and* $C(\mu_0)$ *are mutual relevant positive constants satisfying* $C(\mu_0)$ $\searrow \infty$ *as* $\mu \nearrow 0$.

$$d_{x_0}(x) \;=\; 1 + \frac{|x-x_0|^2}{d_0^2} \quad \text{and} \quad d_0 \;=\; \mathrm{diam}(\Omega).$$

Then our WHHFs are smooth on a neighbourhood of $z_0 = (t_0, x_0)$.

Theorem 1.3 (Second Regularity Theorem). *If the range of* u_0: $u_0\ (\overline{\Omega})$ *is compactly contained in* $\mathbb{S}^D_+$ *aftermath of a suitable rotation if necessary, then the range of our heat flow* $u(\overline{Q}(T))$ *is done there and they are smooth in* $(0,T) \times \overline{\Omega}$.

We list a glossary of notation below: We adopt it from K.Horihata [9] and add it up the new symbols especially related to the boundary. The symbols of vector and the definition on the function class can be seen therein.

Notation

(i) $\mathbb{R}^d_\pm = \{x = (x', x_d) \in \mathbb{R}^d \,;\ x_d \gtrless 0\}$ and
$\mathbb{R}^d_0 = \{x = (x', x_d) \in \mathbb{R}^d \,;\ x_d \,=\, 0\}$ $(\backsimeq \mathbb{R}^{d-1})$.

(ii) For any set $A \subset \mathbb{R}^d$ and any positive number δ, $A_\delta = \{x \in \mathbb{R}^d \,;\, \inf_{y\in\Omega} |x - y| \,<\, \delta\}$.

(iii) The symbol Ω is a domain in $\mathbb{R}^d$ and $\partial\Omega$ the boundary of Ω. Moreover for a set $A \subset\mathbb{R}^d$, the symbol $A_\pm$ respectively means $A \cap \Omega$ and $A \cap \complement\Omega$ and A_0 does $A \cap \partial\Omega$.

(iv) A vector ν and τ respectively denotes the unit outer normal and the tangential field along $\partial\Omega$.

(v) $Q(T) = (0,T) \times \Omega$. $\partial Q(T) \;=\; [0,T) \times \partial\Omega \cup \{0\} \times \Omega$.

(vi) $\mathbb{S}^D \;=\; \{y = (y^1, y^2, \ldots, y^{D+1}) \,\in\, \mathbb{R}^{D+1};\ |y| \;=\; \sqrt{\sum_{i=1}^{D+1} (y^i)^2} \;=\; 1\}$.

(vii) $\mathbb{S}^D_\pm \;=\; \{y = (y', y^{D+1}) = (y^1, y^2, \ldots, y^{D+1}) \,\in\, \mathbb{S}^D;\ y^{D+1} \,\gtrless\, 0\}$.

(viii) $\mathbb{S}^D_0 \;=\; \{y = (y', y^{D+1}) = (y^1, y^2, \ldots, y^{D+1}) \,\in\, \mathbb{S}^D;\ y^{D+1} \,=\, 0\}$
$(\backsimeq \partial B^D_1(0))$.

(ix) Set points $x = (x_1, x_2, \dots, x_d)$, $x_0 = (x_{0,1}, x_{0,2}, \dots, x_{0,d})$ and $z_0 = (t_0, x_0) \in \mathbb{R}^{d+1}$; Then indicate a ball $B_r(x_0)$, a parabolic cylinder $P_r(z_0)$ by

$$\begin{aligned} B_r(x_0) &= \{x \in \mathbb{R}^d\,;\, |x - x_0| < r\}, \\ P_r(z_0) &= \{z = (t, x) \in Q(T)\,;\, t_0 - r^2 < t < t_0 + r^2, |x - x_0| < r\}. \end{aligned}$$

In $B_r(x_0)$ and $P_r(z_0)$, the points of x_0 and z_0 will be often abbreviated when no confusion may arise.

(x) Letter C denotes a generic constant. By the letter $C(B)$, it means that a constant depends only on a parameter B.

(xi) $\kappa(t)$ is $\arctan(t)/\pi$.

(xii) For any point $z_0 = (t_0, x_0)$ in $\mathbb{R}^{d+1}$, the back-ward heat kernel G_{z_0} designates

$$G_{z_0}(t, x) = \frac{1}{\sqrt{4\pi(t_0 - t)}^{\,d}} \exp\Big(-\frac{|x - x_0|^2}{4(t_0 - t)}\Big) \quad ((t, x) \in (0, t_0) \times \mathbb{R}^d).$$

(xiii) For the nabla: $\nabla = \big(\partial/\partial y_1, \partial/\partial y_2, \dots, \partial/\partial y_d\big)$, the differential operators ∇_ν and ∇_τ denote

$$\nabla_\nu = \nu \cdot \nabla, \ \nabla_\tau = \nabla - \nu\nu \cdot \nabla.$$

A forthcoming paper extends the results here and the ones of K.Horihata [9] to a heat flow between any Riemannian manifold under a certain assumption.

2 GLHF.

The chapter introduces three fundamental energy inequalities: The first is a monotonical inequality and the second is a local decay energy estimate, which is can be regarded as a variant of a monotonical inequality, and the final is a hybrid inequality.

2.1 Two Monotonical Inequalities.

We introduce former two inequalities. The one is it by Y.Chen and F.H.Lin [3] and the another is a hybrid type inequality combined with the energy decay estimate referred to K.Horihata [9, Theorem 2.6] and the usual monotonical inequality:

Theorem 2.1 (Monotonical Inequalities). *There exist constants C and $C(\mu_0)$ with $C(\mu_0) \nearrow \infty$ as $\mu_0 \searrow 0$ such that the following holds for any point $z_0 = (t_0, x_0) \in Q(T)$ and positive numbers R, R_1 and R_2 with $0 < R_1 < R_2$ with $\max(R_2, R) \leq \sqrt{t_0}/2$,*

$$\int_{t_0-4R_1^2}^{t_0-R_1^2} dt \int_{\Omega} \mathbf{e}_\lambda \, G_{z_0} \, dx + \int_{R_1}^{R_2} dt \int_{\Omega} \left| \frac{\partial u_\lambda}{\partial t} - \frac{x - x_0}{2(t - t_0)} \cdot \nabla u_\lambda \right|^2 G_{z_0} \, dx$$
$$\leq C \exp\left(R_2^{\mu_0} - R_1^{\mu_0}\right) \int_{t_0-4R_2^2}^{t_0-R_2^2} dt \int_{\Omega} \mathbf{e}_\lambda \, G_{z_0} \, dx$$
$$+ C(\mu_0)(R_2 - R_1) \tag{2.1}$$

and in addition under the hypothesis (B),

$$R^2 \int_{\{t_0-R^2\}\times\Omega} \mathbf{e}_\lambda G_{(t_0-R^2, x_0)} \, dx \leq \frac{e^{-4(d-2)(t_0-R^2)/d_0^2}}{(2\sqrt{t_0})^d} \int_{\Omega} |\nabla u_0|^2 \, dx$$
$$+ C \int_0^{t_0-R^2} dt \int_{\partial\Omega} |\nabla_\tau u_0|^2 \, G_{z_0} \, d\mathcal{H}_x^{d-1} \tag{2.2}$$

with

$$G_{z_0}(t, x) = \frac{1}{\sqrt{4\pi(t_0 - t)}^{\,d}} \exp\Big(-\frac{|x - x_0|^2}{4(t_0 - t)}\Big) \quad ((t, x) \in (0, t_0) \times \mathbb{R}^d),$$
$$d_{x_0}(x) = 1 + \frac{|x - x_0|^2}{d_0^2} \quad \text{with} \quad d_0 = \operatorname{diam}(\Omega).$$

Proof of Theorem 2.1.

Set

$$\nabla_{|x-x_0|} = \frac{x - x_0}{|x - x_0|} \cdot \nabla, \ \nabla_{\tau_0} = \nabla - \frac{x - x_0}{|x - x_0|} \nabla_{|x-x_0|}. \tag{2.3}$$

Since (2.1) is well-known, we only show (2.2): Multiplying (1.8) by

$$2(t_0 - t)\frac{\partial u_\lambda}{\partial t} G_{z_0} d_{x_0} \quad \text{and} \quad (x - x_0) \cdot \nabla u_\lambda G_{z_0} (d_{x_0} + 4(t_0 - t)/d_0^2)$$

and integrating it over Ω, we obtain

$$\begin{aligned}
&2\int\limits_{\Omega}\left|\frac{\partial u_\lambda}{\partial t}\right|^2 (t_0-t)G_{z_0}d_{x_0}\,dx \;+\; 2\frac{d}{dt}\int\limits_{\Omega}\mathbf{e}_\lambda(t_0-t)\,G_{z_0}d_{x_0}\,dx\\
&-\;(d-2)\int\limits_{\Omega}\mathbf{e}_\lambda\,G_{z_0}d_{x_0}\,dx \;+\; \int\limits_{\Omega}\mathbf{e}_\lambda\frac{|x-x_0|^2}{2(t_0-t)}\,G_{z_0}d_{x_0}\,dx\\
&+\;\lambda^{1-\kappa}\int\limits_{\Omega}(|u_\lambda|^2\;-\;1)^2\,\frac{|x-x_0|^2}{8(t_0-t)}\,G_{z_0}\,d_{x_0}\,dx\\
&-\;\int\limits_{\Omega}\Big\langle\frac{\partial u_\lambda}{\partial t},(x-x_0)\cdot\nabla u_\lambda\Big\rangle\,G_{z_0}\big(d_{x_0}\;-\;\frac{4(t_0-t)}{d_0^2}\big)\,dx \qquad\qquad (2.4)\\
&+\;\dot{\kappa}\log\lambda\,\frac{\lambda^{1-\kappa}}{2}\int\limits_{\Omega}(|u_\lambda|^2\;-\;1)^2\,(t_0-t)\,G_{z_0}\,d_{x_0}\,dx\;=\;0,
\end{aligned}$$

$$\begin{aligned}
&-\;\int\limits_{\Omega}\Big\langle\frac{\partial u_\lambda}{\partial t},(x-x_0)\cdot\nabla u_\lambda,\Big\rangle\,G_{z_0}\left(d_{x_0}\;+\;\frac{4(t_0-t)}{d_0^2}\right)dx\\
&+\;(d-2)\int\limits_{\Omega}\mathbf{e}_\lambda\,G_{z_0}\Big(d_{x_0}\;+\;\frac{4(t_0-t)}{d_0^2}\Big)\,dx\\
&+\;\int\limits_{\Omega}(|\nabla_{|x-x_0|}u_\lambda|^2\;-\;|\nabla_{\tau_0}u_0|^2)\,\frac{|x-x_0|^2}{4(t_0-t)}\,G_{z_0}\,d_{x_0}\,dx \qquad\qquad (2.5)\\
&+\;\frac{\lambda^{1-\kappa}}{2}\int\limits_{\Omega}(|u_\lambda|^2\;-\;1)^2\,\frac{|x-x_0|^2}{t_0-t}G_{z_0}\,\big(d_{x_0}\;+\;\frac{4(t_0-t)}{d_0^2}\big)\,dx\\
&-\;\frac{\lambda^{1-\kappa}}{8}\int\limits_{\Omega}(|u_\lambda|^2\;-\;1)^2\,\frac{|x-x_0|^2}{t_0-t}G_{z_0}\,d_{x_0}\,dx\\
&+\;\frac{1}{2}\int\limits_{\partial\Omega}(\langle x-x_0,\nu\rangle|\nabla_\nu u_\lambda|^2\;+\;2\langle\nabla_\nu u_\lambda,(x-x_0)\cdot\nabla_\tau u_0\rangle\;-\;\langle x-x_0,\nu\rangle|\nabla_\tau u_0|^2)\\
&\qquad\times G_{z_0}\left(d_{x_0}\;+\;\frac{4(t_0-t)}{d_0^2}\right)d\mathcal{H}_x^{d-1}\;=\;0.
\end{aligned}$$

Summarizing (2.4), (2.5) and using (B), we arrive at

$$\begin{aligned}
&\frac{d}{dt}\int\limits_{\Omega}\mathbf{e}_\lambda\,(t_0-t)d_{x_0}G_{z_0}\,dx\;+\;\frac{4(d-2)}{d_0^2}\int\limits_{\Omega}\mathbf{e}_\lambda\,(t_0-t)G_{z_0}d_{x_0}\,dx\\
&+\;2\int\limits_{\Omega}\left|\frac{\partial u_\lambda}{\partial t}\sqrt{t_0-t}\;-\;\frac{x-x_0}{2\sqrt{t_0-t}}\cdot\nabla u_\lambda\right|^2G_{z_0}d_{x_0}\,dx\;+\;\frac{\theta_0}{2}\int\limits_{\Omega}|x'\;-\;x_0'|^2|\nabla_\nu u_\lambda|^2G_{z_0}\,d_{x_0}\,dx
\end{aligned}$$

$$\leq C \int\limits_{\partial\Omega} |\nabla_\tau u_0|^2 G_{z_0} \Big(d_{x_0} + \frac{4(t_0 - t)}{d_0^2} \Big) \, d\mathcal{H}_x^{d-1}. \tag{2.6}$$

A multiplier of (2.6) by $e^{4(d-2)t/d_0^2}$ and an integration on it from 0 to $t_0 - R^2$ imply our claim.

2.2 Hybrid type Inequality for GLHF.

We demonstrate the reverse Poincaré inequality. To this end we prepare the below: Let the mapping h_0 be the solution to

$$\begin{cases} -\triangle h_0 &= 0 \quad \text{in} \quad \Omega, \\ \quad h_0 &= u_0 \quad \text{on} \quad \partial\Omega. \end{cases} \tag{2.7}$$

Then we claim

Theorem 2.2 (Hybrid Inequality). *For any positive number ϵ_0 and any point z_0 in $Q(T)$, there exists a positive constant $C(\epsilon_0)$ satisfying $C(\epsilon_0) \nearrow \infty$ as $\epsilon_0 \searrow 0$ such that the inequality*

$$\int\limits_{P_R(z_0)\cap Q(T)} \mathbf{e}_\lambda(z) \, dz \leq \epsilon_0 \int\limits_{P_{2R}(z_0)\cap Q(T)} \mathbf{e}_\lambda(z)\, dz + \frac{C(\epsilon_0)}{R^2} \int\limits_{P_{2R}(z_0)\cap Q(T)} |u_\lambda(z) - h_0(x)|^2 \, dz$$
$$+ \int\limits_{P_{2R}(z_0)\cap Q(T)} (|\nabla^{[(d+1)/2]+1} h_0(x)|^2 \, dz + |\nabla h_0(x)|^2) \, dz + o(1) \quad (\lambda \nearrow \infty) \tag{2.8}$$

holds for any parabolic cylinder $P_{2R}(z_0)$.

Preliminary we rewrite (1.8) to it by the flatten-out coordinate: Set $v_\lambda(t,y)$ on $\Phi_{x_0}(U(x_0)\cap\Omega)$ by $v_\lambda(t,y) = u_\lambda(t, \Phi_{x_0}^{-1}(y))$. Then (1.8) becomes

$$\frac{\partial v_\lambda}{\partial t} - L(v_\lambda) + \lambda^{1-\kappa}\big(|v_\lambda|^2 - 1\big) v_\lambda = 0 \text{ in } (0,T) \times \Phi_{x_0}(U(x_0)\cap\Omega) \tag{2.9}$$

with

$$D_i = \frac{\partial}{\partial y_i} - a_i \frac{\partial}{\partial y_d}, \quad a_i = \frac{\partial \phi_{x_0}}{\partial y_i} \quad (i = 1, 2, \dots, d-1),$$
$$D_d = \frac{\partial}{\partial y_d}, \; L = \sum_{i=1}^{d-1} D_i^2 + D_d^2,$$

$$
\begin{aligned}
D_\nu \;&=\; \sum_{i=1}^{d} \frac{y^i}{|y|} D_i, \;\; D_\tau \;=\; D \;-\; \frac{y}{|y|}\, D_\nu, \\
\triangle_\tau \;&=\; \triangle \;-\; \frac{1}{\rho^{d-1}} \frac{\partial}{\partial \rho}\Big(\rho^{d-1} \frac{\partial}{\partial \rho}\Big).
\end{aligned} \tag{2.10}
$$

Choose $B_R(0) \subset \Phi_{x_0}(U(x_0))$ and induce the mapping h_{x_0} by

$$
h_{x_0}(t, y) \;=\; h_0(\Phi_{x_0}^{-1}(y)). \tag{2.11}
$$

We extend the mapping $v_\lambda \,-\, h_{x_0}$ in $B_R(0) \cap \mathbb{R}^d_+$ to the one in $B_R(0)$ by

$$
\begin{cases}
\;\;(v_\lambda \,-\, h_{x_0})(t, (y', y_d)) & \text{if} \;\; y \,\in\, B_R(0) \cap \mathbb{R}^d_+, \\
\qquad\qquad\qquad\qquad 0 & \text{if} \;\; y \,\in\, B_R(0) \cap \mathbb{R}^d_0, \\
-(v_\lambda \,-\, h_{x_0})(t, (y', -y_d)) & \text{if} \;\; y \,\in\, B_R(0) \cap \mathbb{R}^d_-
\end{cases}
$$

and denote it by the same symbol $v_\lambda - h_{x_0}$: Note that the mapping $v_\lambda - h_{x_0}$ belongs to $(C^1 \cap W^{2,2})\;(B_R(0)\,;\,\mathbb{R}^{D+1})$ and it satisfies the following identity:

$$
\begin{aligned}
&\int_{B_R(0)} \Big\langle \frac{\partial v_\lambda}{\partial t}, \phi \Big\rangle \, dx \;+\; \int_{B_R(0)} \langle D(v_\lambda - h_{x_0}), D\phi \rangle \, dx \\
&\qquad +\, \lambda^{1-\kappa} \int_{B_R(0)} \big(|v_\lambda|^2 \,-\, 1\big) \langle v_\lambda, \phi \rangle \, dx \;=\; 0
\end{aligned} \tag{2.12}
$$

for any mapping $\phi \in C_0^\infty\;(B_R(0)\,;\,\mathbb{R}^{D+1})$ with

$$
\begin{aligned}
\langle D(v_\lambda - h_{x_0}), D\phi \rangle \;&=\; \sum_{i=1}^{d} \sum_{j=1}^{D+1} D_i (v_\lambda - h_{x_0})^j D_i \phi^j \\
\Big\langle \frac{\partial v_\lambda}{\partial t}, \phi \Big\rangle \;&=\; \sum_{j=1}^{D+1} \frac{\partial v_\lambda^j}{\partial t} \phi^j, \;\; \langle v_\lambda, \phi \rangle \;=\; \sum_{j=1}^{D+1} v_\lambda^j \phi^j
\end{aligned}
$$

to verify

Lemma 2.3 *For any balls B_{ρ_1} and B_{ρ_2} with $B_{\rho_1} \,\subset\, B_{\rho_2} \,\subset\, B_R(0)$ and any cylinders P_{ρ_1} and P_{ρ_2} with $P_{\rho_1} \,\subset\, P_{\rho_2} \,\subset\, P_R(0)$, the following holds*:

$$
\begin{aligned}
&\int_{P_{\rho_1}} \lambda^{1-\kappa} (1 \,-\, |v_\lambda|^2) \, dz \;\le\; C \int_{P_{\rho_2}} \mathbf{e}_\lambda \, dz \\
&\qquad +\, \frac{C}{(\rho_2 - \rho_1)^2} \int_{P_{\rho_2}} (1 \,-\, |v_\lambda|^2) \, dz,
\end{aligned} \tag{2.13}
$$

$$\lambda^{1-\kappa}\int\limits_{B_{\rho_1}}(1-|v_\lambda|^2)\,dy \le C\int\limits_{B_{\rho_2}}\mathbf{e}_\lambda\,dy$$
$$+\frac{C}{\rho_2-\rho_1}\int\limits_{B_{\rho_2}}|Dv_\lambda|\,dy + C\int\limits_{B_{\rho_2}}\Big|\frac{\partial v_\lambda}{\partial t}\Big|\,dy. \tag{2.14}$$

Secondly we list symbols and auxiliary mapping employed only here. Give L_λ by $[\log(\lambda/h(\lambda))/\log 2]+1$, where a function $h(\lambda)$ $(\lambda\in\mathbb{R})$ is positive satisfying $h(\lambda)\searrow 0$ $(\lambda\nearrow\infty)$. We then introduce the decompositional convention: Let r be a positive number less than $\epsilon_0^2 R$, and put

$$\triangle\rho_l = C_1\epsilon_0^4 r(1/2)^l \quad (l=1,2,\dots,L_\lambda),$$
$$\rho_l = \begin{cases} 0 & (l=0)\\ (1-\epsilon_0^4)r + C_1\epsilon_0^4 r\sum_{j=1}^{l}(1/2)^j & (l=1,2,\dots,L_\lambda)\end{cases}$$
$$\text{with } C_1 = \Big(\sum_{l=1}^{L_\lambda}(1/2)^l\Big)^{-1}.$$

Fix $t\in(-r^2,r^2)$ and choose numbers r_l $(l=1,2,\dots,L_\lambda-1)$ and $\triangle r_l$ $(l=1,2,\dots,L_\lambda)$ so that they are given by

$$\frac{\triangle\rho_l}{3}\int\limits_{\{r_l\}\times\mathbb{S}^{d-1}}|D_\nu(v_\lambda-h_{x_0})(t,y)|^2\,d\mathcal{H}^{d-1}_y = \int\limits_{\rho_l-\triangle\rho_l/6}^{\rho_l+\triangle\rho_l/6}\rho^{d-1}\,d\rho\int\limits_{\mathbb{S}^{d-1}}|D_\nu(v_\lambda-h_{x_0})(t,y)|^2\,d\omega_{d-1},$$

$r_0=0,\ r_{L_\lambda}=\rho_{L_\lambda}=r, \triangle r_l = r_l - r_{l-1}$.

In addition a positive number $\hat{r}_l$ $(l=0,1,\dots,L_\lambda)$ denotes

$$\widehat{r_l} = \begin{cases} 0 & (l=0)\\ \dfrac{r_l+r_{l+1}}{2} & (l=1,2,\dots,L_\lambda-1)\\ r & (l=L_\lambda).\end{cases}$$

Next introduce a mapping f_λ which is the solution to

$$\begin{cases} D_\nu f_\lambda + r\triangle_\tau f_\lambda = 0 & \text{in } (0,r)\times\mathbb{S}^{d-1},\\ f_\lambda = v_\lambda - h_{x_0} & \text{on } \{r\}\times\mathbb{S}^{d-1}.\end{cases} \tag{2.15}$$

Designate five sorts of annuls:

$$\widehat{T_l} = [\widehat{r}_{l-1},\widehat{r_l})\times\mathbb{S}^{d-1},\quad T_l = [r_{l-1},r_l)\times\mathbb{S}^{d-1},$$

$$
\begin{aligned}
T_l^1 \;&=\; [r_{l-1}, r_{l-1} \,+\, \triangle r_l/3) \,\times\, \mathbb{S}^{d-1},\\
T_l^2 \;&=\; [r_{l-1} \,+\, \triangle r_l/3, r_l \,-\, \triangle r_l/3) \,\times\, \mathbb{S}^{d-1},\\
T_l^3 \;&=\; [r_l \,-\, \triangle r_l/3, r_l) \,\times\, \mathbb{S}^{d-1} \quad (l = 1, 2, \ldots, L_\lambda).
\end{aligned}
\tag{2.16}
$$

A next step establishing a Hybrid type inequality is to construct a certain support mappings $\widetilde{w}_{\lambda,l}$, $\widehat{w}_{\lambda,l}$ and $w_{\lambda,l}$ by making the best of the function f_λ: They are given by the solutions of

$$
\left\{
\begin{array}{rcl}
-\triangle \widetilde{w}_{\lambda,1} &=& 0 \qquad\qquad \text{in} \quad [0, r_1) \times \mathbb{S}^{d-1}\\
\widetilde{w}_{\lambda,1}\big|_{\{r_1\}\times\mathbb{S}^{d-1}} &=& f_\lambda\big|_{\{r_1\}\times\mathbb{S}^{d-1}},
\end{array}
\right.
\tag{2.17}
$$

$$
\left\{
\begin{array}{rcl}
-\triangle \widetilde{w}_{\lambda,l} &=& 0 \qquad \text{in} \quad T_l\\
\widetilde{w}_{\lambda,l}\big|_{\{r_{l-1}\}\times\mathbb{S}^{d-1}} &=& f_\lambda\big|_{\{r_{l-1}\}\times\mathbb{S}^{d-1}} \quad (l = 2, 3, \ldots, L_\lambda),\\
\widetilde{w}_{\lambda,l}\big|_{\{r_l\}\times\mathbb{S}^{d-1}} &=& f_\lambda\big|_{\{r_l\}\times\mathbb{S}^{d-1}}
\end{array}
\right.
\tag{2.18}
$$

$$
\left\{
\begin{array}{rcl}
-\triangle \widehat{w}_{\lambda,1} &=& 0 \qquad\qquad \text{in} \quad [0, \widehat{r}_1) \times \mathbb{S}^{d-1},\\
\widehat{w}_{\lambda,1}\big|_{\{\hat{r}_1\}\times\mathbb{S}^{d-1}} &=& \widetilde{w}_{\lambda,2}\big|_{\{\hat{r}_1\}\times\mathbb{S}^{d-1}}
\end{array}
\right.
\tag{2.19}
$$

$$
\left\{
\begin{array}{rcl}
-\triangle \widehat{w}_{\lambda,l} &=& 0 \qquad \text{in} \quad \widehat{T}_l\\
\widehat{w}_{\lambda,l}\big|_{\{\hat{r}_l\}\times\mathbb{S}^{d-1}} &=& \widetilde{\widetilde{w}}_{\lambda,l+1}\Big|_{\{\hat{r}_l\}\times\mathbb{S}^{d-1}} \quad (l = 2, 3, \ldots, L_\lambda - 1),\\
\widehat{w}_{\lambda,l}\big|_{\{\hat{r}_{l-1}\}\times\mathbb{S}^{d-1}} &=& \widetilde{w}_{\lambda,l}\big|_{\{\hat{r}_{l-1}\}\times\mathbb{S}^{d-1}}
\end{array}
\right.
\tag{2.20}
$$

$$
\left\{
\begin{array}{rcl}
-\triangle \widehat{w}_{\lambda,L_\lambda} &=& 0 \qquad \text{in} \quad \widehat{T}_{L_\lambda}\\
\widehat{w}_{\lambda,L_\lambda}\big|_{\{\hat{r}_{L_\lambda}\}\times\mathbb{S}^{d-1}} &=& u_\lambda\big|_{\{r\}\times\mathbb{S}^{d-1}}\\
\widehat{w}_{\lambda,L_\lambda}\big|_{\{\hat{r}_{L_\lambda-1}\}\times\mathbb{S}^{d-1}} &=& \widetilde{w}_{\lambda,L_\lambda}\big|_{\{\hat{r}_{L_\lambda-1}\}\times\mathbb{S}^{d-1}},
\end{array}
\right.
\tag{2.21}
$$

$$
\left\{ \quad w_{\lambda,1} \;=\; \widehat{w}_{\lambda,1} \qquad \text{on} \quad T_1, \right.
\tag{2.22}
$$

$$
\left\{
\begin{array}{rcl}
-\triangle w_{\lambda,l} &=& 0 \qquad \text{in} \quad T_l\\
w_{\lambda,l}\big|_{\{r_l\}\times\mathbb{S}^{d-1}} &=& \widehat{\widehat{w}}_{\lambda,l}\Big|_{\{r_l\}\times\mathbb{S}^{d-1}} \qquad (l = 2, 3, \ldots, L_\lambda - 1),\\
w_{\lambda,l}\big|_{\{r_{l-1}\}\times\mathbb{S}^{d-1}} &=& \widehat{w}_{\lambda,l-1}\big|_{\{r_{l-1}\}\times\mathbb{S}^{d-1}}
\end{array}
\right.
\tag{2.23}
$$

$$
\left\{
\begin{array}{rcl}
-\triangle w_{\lambda,L_\lambda} &=& 0 \qquad \text{in} \quad T_{L_\lambda}\\
w_{\lambda,L_\lambda}\big|_{\{r_{L_\lambda}\}\times\mathbb{S}^{d-1}} &=& (v_\lambda - h_{x_0})\big|_{\{r\}\times\mathbb{S}^{d-1}}\\
w_{\lambda,L_\lambda}\big|_{\{r_{L_\lambda-1}\}\times\mathbb{S}^{d-1}} &=& \widehat{w}_{\lambda,L_\lambda-1}\big|_{\{r_{L_\lambda-1}\}\times\mathbb{S}^{d-1}},
\end{array}
\right.
\tag{2.24}
$$

where

$$
\begin{aligned}
\widehat{\widehat{w}}_{\lambda,l}(r_l, \omega_{d-1}) \;&= \widehat{w}_{\lambda,l}(r_l, \omega_{d-1})\\
&\qquad -\; (r_l \,-\, \hat{r}_{l-1}) \int\limits_0^1 D_\nu \widehat{w}_{\lambda,l}(\hat{r}_{l-1} + \tau(r_l - \hat{r}_{l-1}), \omega_{d-1})\, d\tau
\end{aligned}
\tag{2.25}
$$

$$- (\hat{r}_{l-1} - r_{l-1}) \int\limits_0^1 D_\nu \widehat{w}_{\lambda,l-1}(r_{l-1} + \tau(\hat{r}_{l-1} - r_{l-1}), \omega_{d-1})\, d\tau,$$

$$\begin{aligned} \widetilde{\widetilde{w}}_{\lambda,l}(\hat{r}_l, \omega_{d-1}) \ &= \widetilde{w}_{\lambda,l+1}(\hat{r}_l, \omega_{d-1}) \\ &\quad - (\hat{r}_l - r_l) \int\limits_0^1 D_\nu \widetilde{w}_{\lambda,l+1}(r_l + \tau(\hat{r}_l - r_l), \omega_{d-1})\, d\tau \\ &\quad - (r_l - \hat{r}_{l-1}) \int\limits_0^1 D_\nu \widetilde{w}_{\lambda,l}(\hat{r}_{l-1} + \tau(r_l - \hat{r}_{l-1}), \omega_{d-1})\, d\tau. \end{aligned} \tag{2.26}$$

In the following we set

$$\rho_l^1 \ = \ \hat{r}_{l-1} + \tau(r_l - \hat{r}_{l-1}), \ \rho_l^2 \ = \ r_{l-1} + \tau(\hat{r}_{l-1} - r_{l-1})$$

for any positive number τ in $(0,1)$.

We state a few fine properties for the mappings $w_{\lambda,l}$, $\widehat{w}_{\lambda,l}$ and $\widetilde{w}_{\lambda,l}$ used below: To explain it we shall recall a sequence of hyper spherical-harmonics $\{\phi_n^{(\alpha)}\}$ $(n = 0, 1, \ldots; \alpha = 1, 2, \ldots, N(n))$, where hyper spherical-harmonics $\phi_n^{(1)}$, $\phi_n^{(2)}$, $\ldots$, $\phi_n^{(N(n))}$ are independent components with degree n.

Writing $x = (\rho, \omega_{d-1})$, they signify

$$\begin{aligned} \widetilde{w}_{\lambda,l}(x) \ &= \ \widetilde{w}_{\lambda,l}(\rho, \omega_{d-1}) \\ &= \sum_{n=1}^{\infty} \sum_{\alpha=1}^{N(n)} \widetilde{a}_n^{(\alpha)} \Big(\frac{\rho}{r_l}\Big)^n \phi_n^{(\alpha)}(\omega_{d-1}) \ + \sum_{n=1}^{\infty} \sum_{\alpha=1}^{N(n)} \widetilde{b}_n^{(\alpha)} \Big(\frac{r_{l-1}}{\rho}\Big)^{n+d-2} \phi_n^{(\alpha)}(\omega_{d-1}) \\ &\quad + \widetilde{a}_0^{(1)} \end{aligned}$$

with

$$\begin{aligned} \widetilde{a}_n^{(\alpha)} \ &= \ \frac{f_\lambda^{n,(\alpha)}(t, r_l) \ - \ f_\lambda^{n,(\alpha)}(t, r_{l-1})\tau_l^{n+d-2}}{1 - \tau_l^{2n+d-2}}, \\ \widetilde{b}_n^{(\alpha)} \ &= \ -\frac{f_\lambda^{n,(\alpha)}(t, r_l)\tau_l^n \ - \ f_\lambda^{n,(\alpha)}(t, r_{l-1})}{1 - \tau_l^{2n+d-2}}, \\ \tau_l \ &= \ \frac{r_{l-1}}{r_l}, \\ f_\lambda^{n,(\alpha)}(t, r_k) \ &= \ \fint_{\mathbb{S}^{d-1}} \langle f_\lambda(t, r_k, \omega_{d-1}), \phi_n^{(\alpha)}(\omega_{d-1}) \rangle \, d\omega_{d-1} \quad (k = l-1, l), \\ \widehat{w}_{\lambda,l}(x) \ &= \ \widehat{w}_{\lambda,l}(\rho, \omega_{d-1}) \end{aligned}$$

$$
\begin{aligned}
&= \sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)} \widehat{a}_n^{(\alpha)} \Big(\frac{\rho}{\hat{r}_l}\Big)^n \phi_n^{(\alpha)}(\omega_{d-1}) \;+\; \sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)} \widehat{b}_n^{(\alpha)} \Big(\frac{\hat{r}_{l-1}}{\rho}\Big)^{n+d-2} \phi_n^{(\alpha)}(\omega_{d-1}) \\
&+\; \widehat{a}_0^{(1)}
\end{aligned}
$$

with

$$
\begin{aligned}
\widehat{a}_n^{(\alpha)} \;&=\; \frac{\widetilde{\widetilde{w}}^{n,(\alpha)}_{\lambda,l+1}(\hat{r}_l) \;-\; \widetilde{w}^{n,(\alpha)}_{\lambda,l}(\hat{r}_{l-1})\hat{\tau}_l^{n+d-2}}{1-\hat{\tau}_l^{2n+d-2}}, \\
\widehat{b}_n^{(\alpha)} \;&=\; -\frac{\widetilde{\widetilde{w}}^{n,(\alpha)}_{\lambda,l+1}(\hat{r}_l)\hat{\tau}_l^{n} \;-\; \widetilde{w}^{n,(\alpha)}_{\lambda,l}(\hat{r}_{l-1})}{1-\hat{\tau}_l^{2n+d-2}},\; \hat{\tau}_l \;=\; \frac{\hat{r}_{l-1}}{\hat{r}_l}, \\
\widetilde{\widetilde{w}}^{n,(\alpha)}_{\lambda,l+1}(\hat{r}_l) \;&=\; \fint_{\mathbb{S}^{d-1}} \langle \widetilde{\widetilde{w}}_{\lambda,l+1}(\hat{r}_l,\omega_{d-1}), \phi_n^{(\alpha)}(\omega_{d-1})\rangle \, d\omega_{d-1}, \\
\widetilde{w}^{n,(\alpha)}_{\lambda,l}(\hat{r}_l) \;&=\; \fint_{\mathbb{S}^{d-1}} \langle \widetilde{w}_{\lambda,l}(\hat{r}_{l-1},\omega_{d-1}), \phi_n^{(\alpha)}(\omega_{d-1})\rangle \, d\omega_{d-1}, \\
w_{\lambda,l}(x) \;&=\; w_{\lambda,l}(\rho,\omega_{d-1}) \\
&= \sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)} a_n^{(\alpha)} \Big(\frac{\rho}{r_l}\Big)^n \phi_n^{(\alpha)}(\omega_{d-1}) \;+\; \sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)} b_n^{(\alpha)} \Big(\frac{r_{l-1}}{\rho}\Big)^{n+d-2} \phi_n^{(\alpha)}(\omega_{d-1}) \\
&+\; a_0^{(1)}
\end{aligned}
$$

with

$$
\begin{aligned}
a_n^{(\alpha)} \;&=\; \frac{\widehat{\widehat{w}}^{n,(\alpha)}_{\lambda,l}(r_l) \;-\; \widehat{w}^{n,(\alpha)}_{\lambda,l-1}(r_{l-1})\tau_l^{n+d-2}}{1-\tau_l^{2n+d-2}}, \\
b_n^{(\alpha)} \;&=\; -\frac{\widehat{\widehat{w}}^{n,(\alpha)}_{\lambda,l}(r_l)\tau_l^{n} \;-\; \widehat{w}^{n,(\alpha)}_{\lambda,l-1}(r_{l-1})}{1-\tau_l^{2n+d-2}},\; \tau_l \;=\; \frac{r_{l-1}}{r_l}, \\
\widehat{\widehat{w}}^{n,(\alpha)}_{\lambda,l}(r_l) \;&=\; \fint_{\mathbb{S}^{d-1}} \langle \widehat{\widehat{w}}_{\lambda,l}(r_l,\omega_{d-1}), \phi_n^{(\alpha)}(\omega_{d-1})\rangle \, d\omega_{d-1}, \\
\widehat{w}^{n,(\alpha)}_{\lambda,l-1}(r_{l-1}) \;&=\; \fint_{\mathbb{S}^{d-1}} \langle \widehat{w}_{\lambda,l-1}(r_{l-1},\omega_{d-1}), \phi_n^{(\alpha)}(\omega_{d-1})\rangle \, d\omega_{d-1}.
\end{aligned}
$$

We must remark that the 0-degree Fourier coefficients $a_0^{(1)}$, $\widehat{a}_0^{(1)}$ and $\widetilde{a}_0^{(1)}$ are independent of ρ by virtue of $-\triangle w_{\lambda,l} = -\triangle \widehat{w}_{\lambda,l} = -\triangle \widetilde{w}_{\lambda,l} = 0$ and $D_\nu f_\lambda + r \triangle_\tau f_\lambda = 0$: (2.15). We mention four technical lemmas:

Lemma 2.4 (Algebraic Inequalities) *For any positive integer l greater than 1 and less than L_λ, we have*

$$0 \;\le\; \frac{1 - \tau_l^{n+d-2}}{1 - \tau_l^{2n+d-2}} + \frac{n\tau_l^n(1 - \tau_l^{d-2})}{1 - \tau_l^{2n+d-2}} \;\le\; C, \tag{2.27}$$

$$\left|\frac{1 - \tau_{l+1}^{n+d-2}}{1 - \tau_{l+1}^{2n+d-2}} - \tau_l^{n+d-2}\frac{1 - \tau_l^{n+d-2}}{1 - \tau_l^{2n+d-2}}\right| \;\le\; Cn\triangle\tau_l, \tag{2.28}$$

$$\left|\frac{\tau_{l+1}^n(1 - \tau_{l+1}^{d-2})}{1 - \tau_{l+1}^{2n+d-2}} - \frac{\tau_l^{2n+d-2}(1 - \tau_l^{d-2})}{1 - \tau_l^{2n+d-2}}\right| n \;\le\; Cn\triangle\tau_l. \tag{2.29}$$

Lemma 2.5 *For any number l ($l = 2, 3, \ldots, L_\lambda$) and a respective arbitrary point $y = (\rho, \omega_{d-1})$ in T_l for $w_{\lambda,l}$, $\hat{T}_l$ for $\widehat{w}_{\lambda,l}$ and T_l for $\widetilde{w}_{\lambda,l}$, it follows that*

$$D_\nu w_{\lambda,l}(y) \;=\; W_{\lambda,l}^1(y) + W_{\lambda,l}^2(y),$$

where

$$W_{\lambda,l}^1(\rho,\omega_{d-1}) \;=\; \sum_{n=1}^{\infty}\frac{n}{\rho}\sum_{\alpha=1}^{N(n)}\frac{1 - \tau_l^{n+d-2}}{1 - \tau_l^{2n+d-2}}\left(\Big(\frac{\rho}{r_l}\Big)^n - \Big(\frac{r_{l-1}}{\rho}\Big)^{n+d-2}\right)\widehat{w}_{\lambda,l-1}^{n,(\alpha)}(r_{l-1})\phi_n^{(\alpha)}(\omega_{d-1}),$$

$$\begin{aligned}W_{\lambda,l}^2(\rho,\omega_{d-1}) \;=\; &\sum_{n=1}^{\infty}\frac{n}{\rho}\sum_{\alpha=1}^{N(n)}\frac{\tau_l^n(1 - \tau_l^{d-2})}{1 - \tau_l^{2n+d-2}}\Big(\frac{r_{l-1}}{\rho}\Big)^{n+d-2}\widehat{w}_{\lambda,l-1}^{n,(\alpha)}(r_{l-1})\phi_n^{(\alpha)}(\omega_{d-1})\\ &- \frac{d-2}{\rho}\sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)}\frac{1 - \tau_l^n}{1 - \tau_l^{2n+d-2}}\Big(\frac{r_{l-1}}{\rho}\Big)^{n+d-2}\widehat{w}_{\lambda,l-1}^{n,(\alpha)}(r_{l-1})\phi_n^{(\alpha)}(\omega_{d-1}),\end{aligned}$$

$$D_\nu \widehat{w}_{\lambda,l}(y) \;=\; \widehat{W}_{\lambda,l}^1(y) + \widehat{W}_{\lambda,l}^2(y),$$

where

$$\widehat{W}_{\lambda,l}^1(\rho,\omega_{d-1}) \;=\; \sum_{n=1}^{\infty}\frac{n}{\rho}\sum_{\alpha=1}^{N(n)}\frac{1 - \hat{\tau}_l^{n+d-2}}{1 - \hat{\tau}_l^{2n+d-2}}\left(\Big(\frac{\rho}{\hat{r}_l}\Big)^n - \Big(\frac{\hat{r}_{l-1}}{\rho}\Big)^{n+d-2}\right)\widetilde{w}_{\lambda,l}^{n,(\alpha)}(\hat{r}_{l-1})\phi_n^{(\alpha)}(\omega_{d-1}),$$

$$\begin{aligned}\widehat{W}_{\lambda,l}^2(\rho,\omega_{d-1}) \;=\; &\sum_{n=1}^{\infty}\frac{n}{\rho}\sum_{\alpha=1}^{N(n)}\frac{\hat{\tau}_l^n(1 - \hat{\tau}_l^{d-2})}{1 - \hat{\tau}_l^{2n+d-2}}\Big(\frac{\hat{r}_{l-1}}{\rho}\Big)^{n+d-2}\widetilde{w}_{\lambda,l}^{n,(\alpha)}(\hat{r}_{l-1})\phi_n^{(\alpha)}(\omega_{d-1})\\ &- \frac{d-2}{\rho}\sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)}\frac{1 - \hat{\tau}_l^n}{1 - \hat{\tau}_l^{2n+d-2}}\Big(\frac{\hat{r}_{l-1}}{\rho}\Big)^{n+d-2}\widetilde{w}_{\lambda,l}^{n,(\alpha)}(\hat{r}_{l-1})\,\phi_n^{(\alpha)}(\omega_{d-1}),\end{aligned}$$

$$\begin{aligned}&D_\nu \widetilde{w}_{\lambda,l}(y)\\ &\quad= \sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)}\frac{f_\lambda^{n,(\alpha)}(t,r_l) - f_\lambda^{n,(\alpha)}(t,r_{l-1})}{1 - \tau_l^{2n+d-2}}\left(\frac{n}{\rho}\Big(\frac{\rho}{r_l}\Big)^n + \tau_l^n\frac{n+d-2}{\rho}\Big(\frac{r_{l-1}}{\rho}\Big)^{n+d-2}\right)\phi_n^{(\alpha)}(\omega_{d-}\end{aligned}$$

$$
\begin{aligned}
&+\sum_{n=1}^{\infty}\frac{n}{\rho}\sum_{\alpha=1}^{N(n)}\frac{1\,-\,\tau_l^{n+d-2}}{1\,-\,\tau_l^{2n+d-2}}\left(\left(\frac{\rho}{r_l}\right)^{n}\,-\,\left(\frac{r_{l-1}}{\rho}\right)^{n+d-2}\right)f_\lambda^{n,(\alpha)}(t,r_{l-1})\phi_n^{(\alpha)}(\omega_{d-1})\\
&+\sum_{n=1}^{\infty}\frac{n}{\rho}\sum_{\alpha=1}^{N(n)}\frac{\tau_l^{n}(1\,-\,\tau_l^{d-2})}{1\,-\,\tau_l^{2n+d-2}}\left(\frac{r_{l-1}}{\rho}\right)^{n+d-2}f_\lambda^{n,(\alpha)}(t,r_{l-1})\phi_n^{(\alpha)}(\omega_{d-1})\\
&-\frac{d-2}{\rho}\sum_{n=1}^{\infty}\sum_{\alpha=1}^{N(n)}\frac{1\,-\,\tau_l^{n}}{1\,-\,\tau_l^{2n+d-2}}\left(\frac{r_{l-1}}{\rho}\right)^{n+d-2}f_\lambda^{n,(\alpha)}(t,r_{l-1})\phi_n^{(\alpha)}(\omega_{d-1}).
\end{aligned}
$$

Lemma 2.6 *For any positive number ρ, the symbol $S_\rho(\omega_{d-1})$ denotes by a geodesic ball on $\mathbb{S}^{d-1}$ centred at ω_{d-1}; Then for any point $y\in T_l^2$,*

$$
|D_\nu w_{\lambda,l}(y)|^2\;\le\;C\fint_{B_{\triangle r_l/6}(y)}|D_\nu w_{\lambda,l}(y)|^2\,dy \tag{2.30}
$$

and for $y\in T_l^1$ and T_l^3,

$$
\begin{aligned}
|D_\nu w_{\lambda,l}(y)|^2\;\le\;&2\fint_{B_{\triangle r_l/6}(y)}|D_\nu \widehat{w}_{\lambda,l-1}(x)|^2\,dx \qquad\qquad (2.31)\\
&+C\fint_{[r_{l-1},r_{l-1}+\triangle r_l/2)\times S_{\triangle\tau_l/2}(\omega_{d-1})}\left(|D_\nu w_{\lambda,l}(x)|^2\,+\,|D_\nu\widehat{w}_{\lambda,l-1}(x)|^2\right)dx,\\
|D_\nu w_{\lambda,l}(y)|^2\;\le\;&2\fint_{B_{\triangle r_l/6}(y)}|D_\nu \widehat{w}_{\lambda,l}(x)|^2\,dx \qquad\qquad (2.32)\\
&+C\fint_{[r_l-\triangle r_l/2,r_l)\times S_{\triangle\tau_l/2}(\omega_{d-1})}\left(|D_\nu w_{\lambda,l}(x)|^2\,+\,|D_\nu\widehat{w}_{\lambda,l}(x)|^2\right)dx
\end{aligned}
$$

respectively holds. Furthermore (2.30), (2.31) and (2.32) imply

$$
\begin{aligned}
&\sum_{l=2}^{L_\lambda-1}\frac{r^2}{\triangle r_l^2}\int_{T_l}|W_{\lambda,l}^1|^2\,dx\,+\,\sum_{l=2}^{L_\lambda-1}\frac{r^2}{\triangle r_l^2}\int_{\hat{T}_l}|\widehat{W}_{\lambda,l}^1|^2\,dx\\
&\le\;Cr\int_{\partial B_r}|D_\tau v_\lambda|^2\,d\mathcal{H}_x^{d-1}\,+\,o(1)\quad(\lambda\nearrow\infty), \qquad\qquad (2.33)
\end{aligned}
$$

$$
|W_{\lambda,l}^2(y)|\;\le\;\frac{C}{r},\quad|\widehat{W}_{\lambda,l}^2(y)|\;\le\;\frac{C}{r}\quad(l\,=\,1,2,\ldots,L_\lambda) \tag{2.34}
$$

for any respective point $y\in T_l$ and $\hat{T}_l$,

$$\begin{aligned}
&\sum_{l=2}^{L_\lambda-1} \triangle r_l \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} (|W^2_{\lambda,l}|^2 + |\widehat{W}^2_{\lambda,l}|)\, d\mathcal{H}^{d-1}_x \\
&\le \frac{C}{r} \int\limits_{\partial B_r} |v_\lambda - h_{x_0}|^2\, d\mathcal{H}^{d-1}_x, && (2.35)
\end{aligned}$$

$$\begin{aligned}
&\sum_{l=1}^{L_\lambda-1} \frac{1}{\triangle r_l} \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} |D_\nu(w_{\lambda,l+1} - w_{\lambda,l})|^2\, d\mathcal{H}^{d-1}_x && (2.36) \\
&\le Cr \int\limits_{\partial B_r} |D_\tau v_\lambda|^2\, d\mathcal{H}^{d-1}_x + \frac{C}{r} \int\limits_{\partial B_r} |v_\lambda - h_{x_0}|^2\, d\mathcal{H}^{d-1}_x + o(1) \quad (\lambda \nearrow \infty),
\end{aligned}$$

$$\begin{aligned}
&\sum_{l=2}^{L_\lambda} \triangle r_l \int\limits_{\{\rho^1_l\}\times\mathbb{S}^{d-1}} |D_\tau D_\nu \widehat{w}_{\lambda,l}|^2\, d\mathcal{H}^{d-1}_x \\
&\le C\epsilon_0^4 r \int\limits_{\partial B_r} |D(v_\lambda - h_{x_0})|^2\, d\mathcal{H}^{d-1}_x + \frac{C(\epsilon_0)}{r} \int\limits_{\partial B_r} |v_\lambda - h_{x_0}|^2\, d\mathcal{H}^{d-1}_x \\
&+ o(1) \quad (\lambda \nearrow \infty). && (2.37)
\end{aligned}$$

From K.Horihata [9, Lemma 2.3], we deduce (2.37). The final lemma is as follows:

Lemma 2.7 *For any positive number r, we have*

$$\begin{aligned}
&r^2 \sum_{l=2}^{L_\lambda} \int\limits_{T_l} \left(|D^2_\nu \widehat{w}_{\lambda,l}|^2 + |D_\nu D_\tau \widehat{w}_{\lambda,l}|^2 + |D^2_\tau \widehat{w}_{\lambda,l}|^2\right) dy \\
&\le Cr \int\limits_{\partial B_r} |D_\tau(v_\lambda - h_{x_0})|^2\, d\mathcal{H}^{d-1}_y + o(1) \quad (\lambda \nearrow \infty) && (2.38)
\end{aligned}$$

and

$$\begin{aligned}
&r^2 \int\limits_{T_1} \left(|D^2_\nu \widehat{w}_{\lambda,1}|^2 + |D_\nu D_\tau \widehat{w}_{\lambda,1}|^2 + |D^2_\tau \widehat{w}_{\lambda,1}|^2\right) dy \\
&\le \frac{C}{r} \int\limits_{\partial B_r} |v_\lambda - h_{x_0}|^2\, d\mathcal{H}^{d-1}_y. && (2.39)
\end{aligned}$$

After the preparation above, we demonstrate the proof of Theorem 2.2.

Proof of Theorem 2.2.

Take the difference between (1.8) and $\triangle w_{\lambda,l} = 0$ on T_l, multiplying it by $-2y \cdot D((v_\lambda - h_{x_0}) - w_{\lambda,l})$, integrate it on T_l and sum up it for l to verify

$$
\begin{aligned}
&- 2 \sum_{l=1}^{L_\lambda} \int\limits_{T_l} \left\langle \frac{\partial v_\lambda}{\partial t}, y \cdot D((v_\lambda - h_{x_0}) - w_{\lambda,l}) \right\rangle \, dy \\
&+ (d-2) \sum_{l=1}^{L_\lambda} \int\limits_{T_l} |D((v_\lambda - h_{x_0}) - w_{\lambda,l})|^2 \, dy + \frac{d\,\lambda^{1-\kappa}}{2} \sum_{l=1}^{L_\lambda} \int\limits_{T_l} (|v_\lambda|^2 - 1)^2 \, dy \\
&+ 2 \sum_{l=1}^{L_\lambda} \int\limits_{T_l} \langle a \cdot D((v_\lambda - h_{x_0}) - w_{\lambda,l}), D_d((v_\lambda - h_{x_0}) - w_{\lambda,l}) \rangle \, dy \\
&+ 2 \sum_{i,j=1}^{d-1} \sum_{l=1}^{L_\lambda} \int\limits_{T_l} y^j \Big(\frac{\partial a_j}{\partial y_i} - \frac{\partial a_i}{\partial y_j} \Big) \langle D_i((v_\lambda - h_{x_0}) - w_{\lambda,l}), D_d((v_\lambda - h_{x_0}) - w_{\lambda,l}) \rangle \, dy \\
&= 2\lambda^{1-\kappa} \sum_{l=1}^{L_\lambda} \int\limits_{T_l} (|v_\lambda|^2 - 1) \langle v_\lambda, y \cdot D h_{x_0} \rangle \, dy \\
&+ 2\lambda^{1-\kappa} \sum_{l=1}^{L_\lambda} \int\limits_{T_l} (|v_\lambda|^2 - 1) \langle v_\lambda, y \cdot D w_{\lambda,l} \rangle \, dy \\
&- 2 \sum_{l=1}^{L_\lambda} \int\limits_{T_l} \langle (L - \triangle) w_{\lambda,l}, y \cdot D((v_\lambda - h_{x_0}) - w_{\lambda,l}) \rangle \, dy \\
&- \sum_{l=1}^{L_\lambda} \Big(\int\limits_{\{r_l\} \times \mathbb{S}^{d-1}} - \int\limits_{\{r_{l-1}\} \times \mathbb{S}^{d-1}} \Big) \Big(\rho - \frac{y_d a \cdot y}{\sqrt{\rho^2 - y_d^2}} \Big) |D_\nu((v_\lambda - h_{x_0}) - w_{\lambda,l})|^2 \, d\mathcal{H}_y^{d-1} \\
&+ \sum_{l=1}^{L_\lambda} \Big(\int\limits_{\{r_l\} \times \mathbb{S}^{d-1}} - \int\limits_{\{r_{l-1}\} \times \mathbb{S}^{d-1}} \Big) \Big(\rho - \frac{y_d a \cdot y}{\sqrt{\rho^2 - y_d^2}} \Big) |D_\tau((v_\lambda - h_{x_0}) - w_{\lambda,l})|^2 \, d\mathcal{H}_y^{d-1} \\
&+ \sum_{l=1}^{L_\lambda} \Big(\int\limits_{\{r_l\} \times \mathbb{S}^{d-1}} - \int\limits_{\{r_{l-1}\} \times \mathbb{S}^{d-1}} \Big) \\
&\qquad\qquad \frac{\rho y_d}{\sqrt{\rho^2 - y_d^2}} \langle a \cdot D_\tau((v_\lambda - h_{x_0}) - w_{\lambda,l}), D_\nu((v_\lambda - h_{x_0}) - w_{\lambda,l}) \rangle \, d\mathcal{H}_y^{d-1}
\end{aligned}
$$

$$
\begin{aligned}
&+\frac{\lambda^{1-\kappa}}{2}\sum_{l=1}^{L_\lambda}\Big(\int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} - \int\limits_{\{r_{l-1}\}\times\mathbb{S}^{d-1}}\Big)\Big(\rho - \frac{y_d a\cdot y}{\sqrt{\rho^2-y_d^2}}\Big)(|v_\lambda|^2-1)^2\,d\mathcal{H}_y^{d-1}\\
&= \text{(I)} + \text{(II)} + \text{(III)} + \cdots + \text{(VII)} \qquad (2.40)
\end{aligned}
$$

with $a\cdot y = \sum_{i=1}^{d-1} a_i y_i$, $a\cdot D = \sum_{i=1}^{d-1} a_i D_i$.

From now on we shall estimate the each term of the right-hand side in (2.40). Primarily we estimate (I): By using the usual Sobolev inequality, we infer

$$
\begin{aligned}
\text{(I)} \le\ &\epsilon_0^2\lambda^{1-\kappa}\int\limits_{B_r}(1-|v_\lambda|^2)\,dy\\
&+\frac{Cr^2\lambda^{1-\kappa}}{\epsilon_0^2}\int\limits_{B_r}(1-|v_\lambda|^2)\,dy \ \fint\limits_{B_{2r}}(|D^{[d/2]+2}h_{x_0}|^2+|Dh_{x_0}|^2)\,dy. \qquad (2.41)
\end{aligned}
$$

Next we estimate (II): We choose a collection of balls $\{B_{\triangle r_1/12}(y_i)\}$ $(i\in I_1)$ with

$$
B_{r_1}\subset\bigcup_{i\in I_1}B_{\triangle r_1/12}(y_i)\subset B_{\hat r_1}
$$

and for any integer l $(l=2,3,\dots,L_\lambda)$, do three ones of balls $\{B_{\triangle r_l/12}(y_i)\}$ $(i\in I_l^1)$, $\{B_{\triangle r_l/12}(y_i)\}$ $(i\in I_l^2)$ and $\{B_{\triangle r_l/12}(y_i)\}$ $(i\in I_l^3)$

$$
\begin{aligned}
&T_l^1\subset\bigcup_{i\in I_l^1}B_{\triangle r_l/12}(y_i),\quad T_l^2\subset\bigcup_{i\in I_l^2}B_{\triangle r_l/12}(y_i),\quad T_l^3\subset\bigcup_{i\in I_l^3}B_{\triangle r_l/12}(y_i),\\
&\max_{j\in I_l^1\cup I_l^2\cup I_l^3}\operatorname{Card}\{i\in I_l^1\cup I_l^2\cup I_l^3\,;\,B_{\triangle r_l/6}(y_i)\cap B_{\triangle r_l/6}(y_j)\neq\emptyset\}
\end{aligned}
$$

is finite and independent of r and l. From Lemma 2.3, we obtain

$$
\begin{aligned}
\lambda^{1-\kappa}\int\limits_{B_{\triangle r_l/12}(y_i)}(1-|v_\lambda|^2)\,dy \le\ & C\int\limits_{B_{\triangle r_l/6}(y_i)}\mathbf{e}\,dy+\frac{C}{\triangle r_l}\int\limits_{B_{\triangle r_l/6}(y_i)}|Dv_\lambda|\,dy\\
&+\int\limits_{B_{\triangle r_l/6}(y_i)}\left|\frac{\partial v_\lambda}{\partial t}\right|\,dy.
\end{aligned}
$$

Decompose $D_\nu w_{\lambda,l}$ into $W_{\lambda,l}^1$ and $W_{\lambda,l}^2$ in Lemma 2.5 and recall (2.30), (2.31) and (2.32) to yield

$$
\text{(II)} \le \lambda^{1-\kappa}\sum_{i\in I_1}\int\limits_{B_{\triangle r_1/12}(y_i)}(1-|v_\lambda|^2)\ |y\cdot Dw_{\lambda,1}|\,dy
$$

$$
\begin{aligned}
&+\ \lambda^{1-\kappa}\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^1\cup I_l^2\cup I_l^3}\ \int\limits_{B_{\triangle r_l/12}(y_i)} (1\,-\,|v_\lambda|^2)\ |y\cdot Dw_{\lambda,l}|\,dy \\
&+ \sum_{i_{L_\lambda}\in I^1_{L_\lambda}\cup I^2_{L_\lambda}\cup I^3_{L_\lambda}}\ \int\limits_{B_{\triangle r_{L_\lambda}/12}(y_i)\,\cap\, B_r} (1\,-\,|v_\lambda|^2)\ |y\cdot Dw_{\lambda,L_\lambda}|\,dy \\
&\leq\ r\sum_{i\in I_1}\ \int\limits_{B_{\triangle r_1/6}(y_i)} \mathbf{e}_\lambda\,dy \fint\limits_{B_{\triangle r_l/6}(y_i)} |D_\nu w_{\lambda,1}|\,dy \\
&+\ r\sum_{i\in I_1}\ \int\limits_{B_{\triangle r_1/6}(y_i)} \left|\frac{\partial v_\lambda}{\partial t}\right| \fint\limits_{B_{\triangle r_l/6}(y_i)} |D_\nu w_{\lambda,1}|\,dy \\
&+ \sum_{i\in I_1}\ \int\limits_{B_{\triangle r_1/6}(y_i)} |Dv_\lambda|\,\frac{dy}{\triangle r_1} \fint\limits_{B_{\triangle r_1/6}(y_i)} |D_\nu w_{\lambda,1}|\,dy \\
&+\ Cr\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^1}\ \int\limits_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda\,dy \fint\limits_{B_{\triangle r_l/6}(y_i)} (|W^1_{\lambda,l}|\,+\,|\widehat{W}^1_{\lambda,l-1}|)\,dy \\
&+\ Cr\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^1}\ \int\limits_{B_{\triangle r_l/6}(y_i)} \left|\frac{\partial v_\lambda}{\partial t}\right|dy \fint\limits_{B_{\triangle r_l/6}(y_i)} (|W^1_{\lambda,l}|\,+\,|\widehat{W}^1_{\lambda,l-1}|)\,dy \\
&+\ Cr\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^1}\ \int\limits_{B_{\triangle r_l/6}(y_i)} |Dv_\lambda|\,\frac{dy}{\triangle r_l} \fint\limits_{B_{\triangle r_l/6}(y_i)} (|W^1_{\lambda,l}|\,+\,|\widehat{W}^1_{\lambda,l-1}|)\,dy \\
&+\ Cr\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^2}\ \int\limits_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda\,dy \fint\limits_{B_{\triangle r_l/6}(y_i)} |W^1_{\lambda,l}|\,dy \\
&+\ r\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^2}\ \int\limits_{B_{\triangle r_l/6}(y_i)} \left|\frac{\partial v_\lambda}{\partial t}\right|dy \fint\limits_{B_{\triangle r_l/6}(y_i)} |W^1_{\lambda,l}|\,dy \\
&+\ Cr\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^2}\ \int\limits_{B_{\triangle r_l/6}(y_i)} |Dv_\lambda|\,\frac{dy}{\triangle r_l} \fint\limits_{B_{\triangle r_l/6}(y_i)} |W^1_{\lambda,l}|\,dy \\
&+\ Cr\sum_{l=2}^{L_\lambda-1}\ \sum_{i\in I_l^3}\ \int\limits_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda\,dy \fint\limits_{B_{\triangle r_l/6}(y_i)} (|W^1_{\lambda,l}|\,+\,|\widehat{W}^1_{\lambda,l}|)\,dy
\end{aligned}
$$

$$
\begin{aligned}
&+ r \sum_{l=2}^{L_\lambda - 1} \sum_{i \in I_l^3} \int_{B_{\triangle r_l/6}(y_i)} \left| \frac{\partial v_\lambda}{\partial t} \right| dy \fint_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}| + |\widehat{W}^1_{\lambda,l}| \right) dy \\
&+ Cr \sum_{l=2}^{L_\lambda - 1} \sum_{i \in I_l^3} \int_{B_{\triangle r_l/6}(y_i)} |Dv_\lambda| \, \frac{dy}{\triangle r_l} \fint_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}| + |\widehat{W}^1_{\lambda,l}| \right) dy \\
&+ r\lambda^{1-\kappa} \sum_{l=2}^{L_\lambda - 1} \sum_{i \in I_l^1} \int_{B_{\triangle r_l/6}(y_i)} \left(1 - |v_\lambda|^2\right) dy \fint_{B_{\triangle r_l/6}(y_i)} \left(|W^2_{\lambda,l}| + |\widehat{W}^2_{\lambda,l-1}| \right) dy \\
&+ r\lambda^{1-\kappa} \sum_{l=2}^{L_\lambda - 1} \sum_{i \in I_l^2} \int_{B_{\triangle r_l/6}(y_i)} \left(1 - |v_\lambda|^2\right) dy \fint_{B_{\triangle r_l/6}(y_i)} |W^2_{\lambda,l}| \, dy \\
&+ r\lambda^{1-\kappa} \sum_{l=2}^{L_\lambda - 1} \sum_{i \in I_l^3} \int_{B_{\triangle r_l/6}(y_i)} \left(1 - |v_\lambda|^2\right) dy \fint_{B_{\triangle r_l/6}(y_i)} \left(|W^2_{\lambda,l}| + |\widehat{W}^2_{\lambda,l}| \right) dy \\
&+ r\lambda^{1-\kappa} \sum_{i \in I^1_{L_\lambda} \cup I^2_{L_\lambda} \cup I^3_{L_\lambda}} \int_{B_{\triangle r_{L_\lambda}/12}(y_i) \cap B_r} \left(1 - |v_\lambda|^2\right) |y \cdot Dw_{\lambda,L_\lambda}| \, dy \\
&\le \epsilon_0^2 \sum_{i \in I_1} \int_{B_{\triangle r_1/6}(y_i)} \mathbf{e}_\lambda \, dy + r^2 \sum_{i \in I_1} \int_{B_{\triangle r_1/6}(y_i)} \left| \frac{\partial v_\lambda}{\partial t} \right|^2 dy + \epsilon_0^2 \sum_{i \in I_1} \int_{B_{\triangle r_1/6}(y_i)} |Dv_\lambda|^2 \, dy \\
&+ \frac{C}{\epsilon_0^2} \sum_{i \in I_1} \int_{B_{\triangle r_1/6}(y_i)} \mathbf{e}_\lambda \, dy \fint_{B_{\triangle r_1/6}(y_i)} |y \cdot Dw_{\lambda,1}|^2 \, dy \\
&+ \frac{1}{\triangle r_1^2} \sum_{i \in I_1} \int_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,1}|^2 + |\widehat{W}^1_{\lambda,1}|^2 \right) dy \\
&+ \frac{C}{\epsilon_0^2 \triangle r_1^2} \sum_{i \in I_1} \int_{B_{\triangle r_1/6}(y_i)} |y \cdot Dw_{\lambda,1}|^2 \, dy \\
&+ \frac{C}{\epsilon_0^2} \sum_{l=2}^{L_\lambda - 1} \sum_{i \in I_l^1 \cup I_l^2 \cup I_l^3} \int_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda \, dy + \frac{Cr^2}{\epsilon_0^2} \sum_{l=2}^{L_\lambda - 1} \sum_{i \in I_l^1 \cup I_l^2 \cup I_l^3} \int_{B_{\triangle r_l/6}(y_i)} \left| \frac{\partial v_\lambda}{\partial t} \right|^2 dy \\
&+ \frac{1}{\epsilon_0^2 \triangle r_l^2} \sum_{i \in I_1} \int_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,1}|^2 + |\widehat{W}^1_{\lambda,1}|^2 \right) dy
\end{aligned}
$$

$$
\begin{aligned}
&+ C\epsilon_0^2 r^2 \sum_{l=2}^{L_\lambda-1} \sum_{i\,\in\, I_l^1} \int_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda\,dy \fint_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}|^2 + |W^1_{\lambda,l-1}|^2\right) dy \\
&+ C\epsilon_0^2 r^2 \sum_{l=2}^{L_\lambda-1} \sum_{i\,\in\, I_l^2} \int_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda\,dy \fint_{B_{\triangle r_l/6}(y_i)} |W^1_{\lambda,l}|^2\,dy \\
&+ C\epsilon_0^2 r^2 \sum_{l=2}^{L_\lambda-1} \sum_{i\,\in\, I_l^3} \int_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda\,dy \fint_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}|^2 + |\widehat{W}^1_{\lambda,l}|^2\right) dy \\
&+ C\epsilon_0^2 \sum_{l=2}^{L_\lambda-1} \sum_{i\,\in\, I_l^1} \int_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}|^2 + |\widehat{W}^1_{\lambda,l-1}|^2\right) dy \\
&+ C\epsilon_0^2 \sum_{l=2}^{L_\lambda-1} \sum_{i\,\in\, I_l^2} \int_{B_{\triangle r_l/6}(y_i)} |W^1_{\lambda,l}|^2\,dy \\
&+ C\epsilon_0^2 \sum_{l=2}^{L_\lambda-1} \sum_{i\,\in\, I_l^3} \int_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}|^2 + |\widehat{W}^1_{\lambda,l}|^2\right) dy \\
&+ C\epsilon_0^2 \sum_{l=2}^{L_\lambda-1} \left(\frac{r}{\triangle r_l}\right)^2 \sum_{i\,\in\, I_l^1} \int_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}|^2 + |\widehat{W}^1_{\lambda,l-1}|^2\right) dy \\
&+ C\epsilon_0^2 \sum_{l=2}^{L_\lambda-1} \left(\frac{r}{\triangle r_l}\right)^2 \sum_{i\,\in\, I_l^2} \int_{B_{\triangle r_l/6}(y_i)} |W^1_{\lambda,l}|^2\,dy \\
&+ C\epsilon_0^2 \sum_{l=2}^{L_\lambda-1} \left(\frac{r}{\triangle r_l}\right)^2 \sum_{i\,\in\, I_l^3} \int_{B_{\triangle r_l/6}(y_i)} \left(|W^1_{\lambda,l}|^2 + |\widehat{W}^1_{\lambda,l}|^2\right) dy \\
&+ C\lambda^{1-\kappa} \sum_{l=2}^{L_\lambda-1} \sum_{i\,\in\, I_l^1\cup I_l^2\cup I_l^3} \int_{B_{\triangle r_l/12}(y_i)} \left(1 - |v_\lambda|^2\right) dy \\
&+ \lambda^{1-\kappa} \sum_{i\in I^1_{L_\lambda}\cup I^2_{L_\lambda}\cup I^3_{L_\lambda}} \int_{B_{\triangle r_{L_\lambda}/12}(y_i)\,\cap\, B_r} \left(1 - |v_\lambda|^2\right) |y\cdot Dw_{\lambda,L_\lambda}|\,dy. \qquad (2.42)
\end{aligned}
$$

On account of Theorem 2.1,

$$\begin{aligned}
&\frac{1}{(\triangle r_l/6)^{d-2}} \int_{B_{\triangle r_l/6}(y_i)} \mathbf{e}_\lambda(t,y)\,dy \\
&\le \frac{C}{(\triangle r_l/3)^d} \int_{t-(\triangle r_l/3)^2}^{t} ds \int_{B_{\triangle r_l/3}(y_i)} \mathbf{e}_\lambda\,dy + C\triangle r_l^2 \sup_{B_{\triangle r_l/3}} |Dh_{x_0}|^2 \\
&\le \int_{t-(\triangle r_l/3)^2}^{t} ds \int_{\Omega} \mathbf{e}_\lambda G_{(t+(\triangle r_l/3)^2,x)}\,dy + C\triangle r_l^2 \\
&\le C\int_{\Omega} |Du_0|^2\,dy + C\triangle r_l^d
\end{aligned}$$

follows for any time $t \in (t_0 - R^2, t_0 + R^2)$ with a positive number R less than $\sqrt{t_0}/2$ and any ball $B_{\triangle r_l/6}(y_i)$, according to (2.32), (2.33) and (2.34) of Lemma 2.6, which proceeds to our evaluation as follows:

$$\begin{aligned}
\text{(II)} \;\le\; & C\epsilon_0^2 \int_{B_r} \mathbf{e}_\lambda\,dy + \frac{Cr^2}{\epsilon_0^2}\int_{B_r} \left|\frac{\partial v_\lambda}{\partial t}\right|^2 dy + \lambda^{1-\kappa} \int_{B_r\setminus B_{(1-\epsilon_0^4)r}} (1 - |v_\lambda|^2)\,dy \\
&+ \frac{C}{\epsilon_0^2} \int_{B_r\setminus B_{(1-\epsilon_0^4)r}} \mathbf{e}_\lambda\,dy + C\epsilon_0^2 r \int_{\partial B_r} |D(v_\lambda - h_{x_0})|^2\,d\mathcal{H}_y^{d-1} \\
&+ \frac{C(\epsilon_0)}{r} \int_{\partial B_r} |v_\lambda - h_{x_0}|^2\,d\mathcal{H}_y^{d-1} + o(1) \quad (\lambda \nearrow \infty).
\end{aligned} \tag{2.43}$$

By using Lemma 2.7 we similarly asses (III):

$$\begin{aligned}
\text{(III)} \;\le\; & \frac{\epsilon_0^2}{2}\int_{T_1} |D_\nu((v_\lambda - h_{x_0}) - w_{\lambda,1})|^2\,dy + \frac{r^2}{2\epsilon_0^2}\int_{T_1} |(L-\triangle)w_{\lambda,1}|^2\,dy \\
&+ \frac{1}{2\epsilon_0^2}\sum_{l=2}^{L_\lambda}\int_{T_l} |D_\nu(v_\lambda - h_{x_0})|^2\,dy + \frac{(r\epsilon_0)^2}{2}\sum_{l=2}^{L_\lambda}\int_{T_l} |(L-\triangle)w_{\lambda,l}|^2\,dy \\
\le\; & C\epsilon_0^2 \int_{B_r} |D_\nu(v_\lambda - h_{x_0})|^2\,dy + C\epsilon_0^2 r\int_{\partial B_r} |D_\tau(v_\lambda - h_{x_0})|^2\,d\mathcal{H}_y^{d-1} \\
&+ \frac{C}{\epsilon_0^2}\int_{B_r\setminus B_{(1-\epsilon_0^4)r}} |D_\nu(v_\lambda - h_{x_0})|^2\,dy
\end{aligned}$$

$$+\ \frac{C(\epsilon_0)}{r}\int\limits_{\partial B_r}|v_\lambda\ -\ h_{x_0}|^2\,d\mathcal{H}^{d-1}_y\ +\ o(1)\quad(\lambda\nearrow\infty). \tag{2.44}$$

$$\begin{aligned}
\text{(IV)}\ \le\ & 2\sum_{l=1}^{L_\lambda-1}\ \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}}\big(r_l\ -\ \frac{y_d\,a\cdot y}{\sqrt{r_l^2-y_d^2}}\big)\langle D_\nu(v_\lambda-h_{x_0}),D_\nu(w_{\lambda,l}\ -\ w_{\lambda,l+1})\rangle\,d\mathcal{H}^{d-1}_y\\
&-\sum_{l=1}^{L_\lambda-1}\ \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}}\big(r_l\ -\ \frac{y_d\,a\cdot y}{\sqrt{r_l^2-y_d^2}}\big)\langle D_\nu(w_{\lambda,l}\ +\ w_{\lambda,l+1}),D_\nu(w_{\lambda,l}\ -\ w_{\lambda,l+1})\rangle\,d\mathcal{H}^{d-1}_y\\
&\le\ \epsilon_0^6 r\int\limits_{\{r_1\}\times\mathbb{S}^{d-1}}|D_\nu(v_\lambda-h_{x_0})|^2\,d\mathcal{H}^{d-1}_y\\
&+\ \frac{Cr}{\epsilon_0^6}\int\limits_{\{r_1\}\times\mathbb{S}^{d-1}}(|D_\nu w_{\lambda,1}|^2\ +\ |D_\nu w_{\lambda,2}|^2)\,d\mathcal{H}^{d-1}_y\\
&+\ \frac{C}{\epsilon_0^2}\sum_{l=2}^{L_\lambda-1}\triangle r_l\int\limits_{\{r_l\}\times\mathbb{S}^{d-1}}|D_\nu(v_\lambda-h_{x_0})|^2\,d\mathcal{H}^{d-1}_y\\
&+\ \frac{C}{\epsilon_0^2}\sum_{l=2}^{L_\lambda-1}\triangle r_l\int\limits_{\{r_l\}\times\mathbb{S}^{d-1}}(|D_\nu w_{\lambda,l}|^2\ +\ |D_\nu w_{\lambda,l+1}|^2)\,d\mathcal{H}^{d-1}_y\\
&+\ \epsilon_0^2\sum_{l=2}^{L_\lambda-1}\frac{r^2}{\triangle r_l}\int\limits_{\{r_l\}\times\mathbb{S}^{d-1}}|D_\nu(w_{\lambda,l}\ -\ w_{\lambda,l+1})|^2\,d\mathcal{H}^{d-1}_y.
\end{aligned}$$

Using Lemma 2.6, from a definition on r_l, we arrive at

$$\begin{aligned}
\text{(IV)}\ \le\ & C\epsilon_0^2\int\limits_{B_r}|D(v_\lambda-h_{x_0})|^2\,dy\ +\ \frac{C}{\epsilon_0^2}\int\limits_{B_r\setminus B_{(1-\epsilon_0^4)r}}|D(v_\lambda-h_{x_0})|^2\,dy\\
&+\ C\epsilon_0^2 r\int\limits_{\partial B_r}|D(v_\lambda-h_{x_0})|^2\,d\mathcal{H}^{d-1}_y\ +\ \frac{C(\epsilon_0)}{r}\int\limits_{\partial B_r}|v_\lambda-h_{x_0}|^2\,d\mathcal{H}^{d-1}_y\\
&+\ o(1)\quad(\lambda\nearrow\infty).
\end{aligned} \tag{2.45}$$

From $\widehat{\widehat{w}}_{\lambda,L_\lambda}=(v_\lambda-h_{x_0})$ on $\{r\}\times\mathbb{S}^{d-1}$, we obtain

$$\text{(IV)}\ =\ \sum_{l=1}^{L_\lambda}r_l\int\limits_{\{r_l\}\times\mathbb{S}^{d-1}}|D_\tau((v_\lambda-h_{x_0})\ -\ w_{\lambda,l})|^2\,d\mathcal{H}^{d-1}_y$$

$$
\begin{aligned}
&- \sum_{l=1}^{L_\lambda} r_{l-1} \int\limits_{\{r_{l-1}\}\times\mathbb{S}^{d-1}} |D_\tau((v_\lambda - h_{x_0}) \,-\, w_{\lambda,l})|^2 \, d\mathcal{H}^{d-1}_y \\
&= \sum_{l=1}^{L_\lambda-1} r_l \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} |D_\tau((v_\lambda - h_{x_0}) \,-\, \widehat{\widehat{w}}_{\lambda,l})|^2 \, d\mathcal{H}^{d-1}_y \\
&- \sum_{l=1}^{L_\lambda-1} r_l \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} |D_\tau((v_\lambda - h_{x_0}) \,-\, \widehat{w}_{\lambda,l})|^2 \, d\mathcal{H}^{d-1}_y \\
&= -2 \sum_{l=2}^{L_\lambda-1} r_l \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} \langle D_\tau(v_\lambda - h_{x_0}), D_\tau(\widehat{\widehat{w}}_{\lambda,l} \,-\, \widehat{w}_{\lambda,l})\rangle \, d\mathcal{H}^{d-1}_y \\
&+ \sum_{l=2}^{L_\lambda-1} r_l \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} \langle D_\tau(\widehat{\widehat{w}}_{\lambda,l} \,+\, \widehat{w}_{\lambda,l}), D_\tau(\widehat{\widehat{w}}_{\lambda,l} \,-\, \widehat{w}_{\lambda,l})\rangle \, d\mathcal{H}^{d-1}_y.
\end{aligned}
$$

By recalling definition on $\widehat{\widehat{w}}_{\lambda,l}$, namely using (2.25) and (2.37) in Lemma 2.6 to forego to estimate

$$
\begin{aligned}
\text{(V)} \;\le\; & \frac{C}{\epsilon_0^2} \sum_{l=2}^{L_\lambda} \triangle r_l \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} (|D_\tau(v_\lambda - h_{x_0})|^2 \,+\, |D_\tau \widehat{w}_{\lambda,l}|^2) \, d\mathcal{H}^{d-1}_y \\
&+ C\epsilon_0^2 r^2 \sum_{l=1}^{L_\lambda-1} \triangle r_l \int\limits_{\{r_l\}\times\mathbb{S}^{d-1}} \Big(|D_\tau D_\nu \widehat{w}_{\lambda,l}(\rho_l^1, \omega_{d-1})|^2 \,+\, \Big| D_\tau D_\nu \widehat{w}_{\lambda,l-1}(\rho_l^2, \omega_{d-1}) \Big|^2 \Big) \, d\mathcal{H}^{d-1}_y \\
&\le \frac{C}{\epsilon_0^2} \int\limits_{B_r \setminus B_{(1-\epsilon_0^4)r}} |D(v_\lambda - h_{x_0})|^2 \, dy \,+\, C\epsilon_0^2 r \int\limits_{\partial B_r} |D(v_\lambda - h_{x_0})|^2 \, d\mathcal{H}^{d-1}_y \\
&+ \frac{C(\epsilon_0)}{r} \int\limits_{\partial B_r} |v_\lambda - h_{x_0}|^2 \, d\mathcal{H}^{d-1}_y \,+\, o(1) \quad (\lambda \nearrow \infty). \qquad (2.46)
\end{aligned}
$$

The estimate on (VI) is as same as above. Finally we find that (VII) becomes

$$
\text{(VII)} \;=\; \frac{\lambda^{1-\kappa}}{2} \int\limits_{\partial B_r} \big(r - \frac{y_d \, y \cdot a}{\sqrt{r^2 \,-\, y_d^2}} \big) (|v_\lambda|^2 \,-\, 1)^2 \, d\mathcal{H}^{d-1}_y. \qquad (2.47)
$$

On the other hand, recalling the smoothness of a given in (B) in page 1, we also have the following estimate for the left-hand side in (2.40), which is called (L):

$$
\begin{aligned}
(\mathrm{L}) \ \geq\ & \frac{d-2}{4} \int\limits_{B_{(1-\epsilon_0^4)r}} \mathbf{e}_\lambda \, dy \ - \ C\, r^2 \int\limits_{B_r} \left| \frac{\partial v_\lambda}{\partial t} \right|^2 \, dy \\
& - \ \frac{C(\epsilon_0)}{r} \int\limits_{\partial B_r} |v_\lambda - h_{x_0}|^2 \, d\mathcal{H}_y^{d-1} \ - \ C \int\limits_{B_r} |Dh_{x_0}|^2 \, dx
\end{aligned}
\tag{2.48}
$$

A substitution of (2.41), (2.43), (2.44), (2.45), (2.46), (2.47), (2.48) for (2.40), an integration of it with respect to $t \in (-r^2, r^2)$ verifies

$$
\begin{aligned}
\int\limits_{P_{r/2}} \mathbf{e}_\lambda \, dz \ \leq \ & C\epsilon_0^2 \int\limits_{P_r} \mathbf{e}_\lambda \, dz \ + \ \frac{Cr^2}{\epsilon_0^2} \int\limits_{P_r} \left| \frac{\partial v_\lambda}{\partial t} \right|^2 dz \ + \ \frac{C}{\epsilon_0^2} \int\limits_{-r^2}^{r^2} dt \int\limits_{B_r \setminus B_{(1-\epsilon_0^4)r}} |D(v_\lambda - h_{x_0})|^2 \, dy \\
& + \ C\epsilon_0^2 \int\limits_{P_r} \lambda^{1-\kappa} (1 \ - \ |v_\lambda|^2) \, dz \\
& + \ \frac{C}{\epsilon_0^{10}} \int\limits_{P_r} \lambda^{1-\kappa} (1 \ - \ |v_\lambda|^2) \, dz \Big(\fint\limits_{\partial B_r} |v_\lambda \ - \ h_{x_0}|^2 \, d\mathcal{H}_y^{d-1} \ + \ \fint\limits_{\partial B_r} |h_{x_0} \ - \ a|^2 \, d\mathcal{H}_y^{d-1} \Big) \\
& + \ \int\limits_{-r^2}^{r^2} \lambda^{1-\kappa} \, dt \int\limits_{B_r \setminus B_{(1-\epsilon_0^4)r}} (1 \ - \ |v_\lambda|^2) \, dy \ + \ \frac{C(\epsilon_0)}{r} \int\limits_{-r^2}^{r^2} dt \int\limits_{\partial B_r} |v_\lambda \ - \ h_{x_0}|^2 \, d\mathcal{H}_y^{d-1} \\
& + \ \int\limits_{-r^2}^{r^2} \frac{r\lambda^{1-\kappa}}{2} \, dt \int\limits_{\partial B_r} (|v_\lambda|^2 \ - \ 1)^2 \, d\mathcal{H}_y^{d-1} \ + \ C \int\limits_{P_r} |Dh_{x_0}|^2 \, dz \ + \ o(1) \quad (\lambda \nearrow \infty).
\end{aligned}
\tag{2.49}
$$

Integrate (2.49) from $R/2$ to R with respect to r and divide it by R to obtain

$$
\begin{aligned}
\int\limits_{P_{R/4}} \mathbf{e}_\lambda \, dz \ \leq \ & C\epsilon_0^2 \int\limits_{P_R} \mathbf{e}_\lambda \, dz \ + \ \frac{C}{\epsilon_0^2 R} \int\limits_{-R^2}^{R^2} dt \int\limits_{R/2}^{R} dr \int\limits_{B_r \setminus B_{(1-\epsilon_0^4)r}} |D(v_\lambda - h_{x_0})|^2 \, dy \\
& + \ \frac{1}{R} \int\limits_{-R^2}^{R^2} \lambda^{1-\kappa} \, dt \int\limits_{R/2}^{R} dr \int\limits_{B_r \setminus B_{(1-\epsilon_0^4)r}} (1 \ - \ |v_\lambda|^2) \, dy \\
& + \ \frac{CR^2}{\epsilon_0^2} \int\limits_{P_R} \left| \frac{\partial v_\lambda}{\partial t} \right|^2 dz \ + \ \frac{C(\epsilon_0)}{R^2} \int\limits_{P_R} |v_\lambda - h_{x_0}|^2 \, dz
\end{aligned}
$$

$$\begin{aligned}
&+ \frac{1}{R^{d+2}} \int_{P_R} \lambda^{1-\kappa}(1 - |v_\lambda|^2)\,dz \fint_{P_R} |v_\lambda - h_{x_0}|^2\,dz \\
&+ \frac{1}{R^{d-2}} \int_{P_R} \lambda^{1-\kappa}(1 - |v_\lambda|^2)\,dz \fint_{P_R} \Big|\frac{\partial v_\lambda}{\partial t}\Big|^2\,dz \qquad (2.50) \\
&+ \frac{1}{2} \int_{-R^2}^{R^2} \lambda^{1-\kappa}\,dt \int_{B_R \setminus B_{R/2}} (|v_\lambda|^2 - 1)^2\,dy \\
&+ C \int_{P_R} (|D^{[(d+1)/2]+1} h_{x_0}|^2 + |Dh_{x_0}|^2)\,dz + o(1) \quad (\lambda \nearrow \infty).
\end{aligned}$$

By interchanging the integration of the second and the third terms on the right-hand side in (2.50) and employing Lemma 2.3, we claim

$$\begin{aligned}
\int_{P_{R/4}} \mathbf{e}_\lambda\,dz \;\le\; & C\epsilon_0^2 \int_{P_R} \mathbf{e}_\lambda\,dz + \frac{CR^2}{\epsilon_0^2} \int_{P_R} \Big|\frac{\partial v_\lambda}{\partial t}\Big|^2\,dz \\
&+ \frac{C(\epsilon_0)}{R^2} \int_{P_R} |v_\lambda - h_{x_0}|^2\,dz \\
&+ C \int_{P_R} (|D^{[(d+1)/2]+1} h_{x_0}|^2 + |Dh_{x_0}|^2)\,dz + o(1) \quad (\lambda \nearrow \infty). \qquad (2.51)
\end{aligned}$$

The rest of our proof is perfectly as same as the one by K.Horihata [9].

3 WHHF

3.1 Partial Regularity

This chapter studies a partial boundary regularity on WHHF constructed in K.Horihata [9]. We review two convergent theorems:

Theorem 3.1 (Convergence). *There exist a sub-sequence* $\{u_{\lambda(\nu)}\}$ $(\nu = 1, 2, \ldots)$ *of* $\{u_\lambda\}$ $(\lambda > 0)$ *in* $V_B(Q(T); \mathbb{S}^D)$ *and a mapping* $u \in V_B(Q(T); \mathbb{S}^D)$ *such that the sequence of mappings* $\{u_{\lambda(\nu)}\}$ $(\nu = 1, 2, \ldots)$ *respectively converges weakly and weakly-$*$ to a mapping* u *in* $H^{1,2}(0, T; L^2_B(\Omega; \mathbb{R}^{D+1}))$ *and* $L^\infty(0, T; H^{1,2}_B(\Omega; \mathbb{R}^{D+1}))$. *So does it strongly to the mapping* u *in* $L^2((0, T); L^2_B(\Omega; \mathbb{S}^D))$ *and point-wisely to it in almost all* $z \in Q(T)$ *as* $\nu \nearrow \infty$.

Theorem 3.1 enables us state the following existence theorem:

Theorem 3.2 (Existence). *The GLHF converges to a WHHF in* $L^2((0,T); L^2_B(\Omega; \mathbb{S}^D))$ *as* $\lambda \nearrow \infty$ *(modulo a sub-sequence of* λ*).*

Definition 3.3 *Let* $\{u_{\lambda(\nu)}\}$ $(\nu = 1, 2, \ldots)$ *be the sequence selected above and set* $\mathbf{e}_{\lambda(\nu)}$ *the Ginzburg-Landau energy density* $|\nabla u_{\lambda(\nu)}|^2/2 + \lambda(\nu)^{1-\kappa} (|u_{\lambda(\nu)}|^2 - 1)^2/4$. *We then denotes* $\overline{\mathcal{M}}$ *by*

$$\overline{\mathcal{M}}(P_R(z_0) \cap Q(T)) = \limsup_{\lambda(\nu) \nearrow \infty} \frac{1}{R^d} \int\limits_{P_R(z_0) \cap Q(T)} \mathbf{e}_{\lambda(\nu)} \, dz,$$

where $P_R(z_0)$ *is an arbitrary parabolic cylinder.*

Lemma 3.4 (Measured Hybrid Inequality). *Assume that a sequence of GLHF* $\{u_{\lambda(\nu)}\}$ $(\nu = 1, 2, \ldots)$, *respectively converges weakly and weakly-∗ in* $H^{1,2}(0, T; L^2_B(\mathbb{B}^d; \mathbb{R}^{D+1}))$ *and* $L^\infty(0, T; H^{1,2}_B(\mathbb{B}^d; \mathbb{R}^{D+1}))$ *to a WHHF* $u \in V_B(Q; \mathbb{S}^D)$ *as* $\lambda(\nu) \nearrow \infty$. *Then take the pass to the limit* $\lambda(\nu) \nearrow \infty$ *in Theorem 2.2 to infer the following: For any positive number* ϵ_0, *there exists a positive constant* $C(\epsilon_0)$ *satisfying* $C(\epsilon_0) \nearrow \infty$ *as* $\epsilon_0 \searrow 0$ *such that the inequality*

$$\begin{aligned} \overline{\mathcal{M}}\big(P_R(z_0) \cap Q(T)\big) &\le \epsilon_0 \overline{\mathcal{M}}\big(P_{2R}(z_0) \cap Q(T)\big) \\ &+ C(\epsilon_0) \fint\limits_{P_{2R}(z_0) \cap Q(T)} |u(z) - h_0(x)|^2 \, dz \\ &+ C \fint\limits_{P_{2R} \cap Q(T)} (|\nabla^{[(d+1)/2]+1} h_0(x)|^2 + |\nabla h_0(x)|^2) \, dz \end{aligned} \tag{3.1}$$

holds for any parabolic cylinder $P_{2R}(z_0)$.

Likewise L.Simon [15, Lemma 2, p.31], we can assert the following reverse Poincaré inequality.

Corollary 3.5 (Reverse Poincaré Inequality). *The inequality* (3.1) *implies that the following*

$$\begin{aligned} \overline{\mathcal{M}}(P_R(z_0) \cap Q(T)) &\le C \fint\limits_{P_{2R}(z_0) \cap Q(T)} |u(z) - h_0(x)|^2 \, dz \\ &+ C \fint\limits_{P_{2R} \cap Q(T)} (|\nabla^{[(d+1)/2]+1} h_0(x)|^2 + |\nabla h_0(x)|^2) \, dz \end{aligned} \tag{3.2}$$

holds whenever $P_{2R}(z_0)$ *is an arbitrary parabolic cylinder.*

By combining Corollary 3.5 with passing to the limit $\lambda \nearrow \infty$ with Sobolev imbedding theorem and Poincaré inequality for the space variables, we can describe the following lemma. We refer the proof to Theorem 2.1 in M.Giaquinta and M.Struwe [7].

Lemma 3.6 *There exists a positive number* q_0 *greater than* 1 *such that differentials* ∇u *of the WHHF* u *belongs to* $L^{2q_0}((0,T); L_B^{2q_0}(\Omega; \mathbb{R}^{d(D+1)}))$ *with*

$$\begin{aligned}\Big(\fint_{P_R(z_0)} |\nabla u(z)|^{2q_0}\, dz\Big)^{1/2q_0} &\le C\Big(\fint_{P_{2R}(z_0)} |\nabla u(z)|^2\, dz\Big)^{1/2}\\ + C\Big(\fint_{P_{2R}(z_0)} &(|\nabla^{[(d+1)/2]+1} h_0(x)|^2 + |\nabla h_0(x)|^2)\, dz\Big)^{1/2q_0} \qquad (3.3)\end{aligned}$$

for any parabolic cylinder P_{2R}.

If we apply (3.2) in Corollary 3.5 to the proof on a partial regularity result by Y.Chen [2, Lemma 3.1], we can claim

Theorem 3.7 *For any positive number* ϵ, *set*

$$\mathbf{sing}(\epsilon) = \bigcap_{R>0} \{z_0 \in (0,T)\times\overline{\Omega};\ \overline{\mathcal{M}}\big(P_R(z_0)\big) \ge \epsilon\}, \qquad (3.4)$$

$$\mathbf{reg}(\epsilon) = (0,T)\times\overline{\Omega} \setminus \mathbf{sing}. \qquad (3.5)$$

Then there exist some positive number ϵ_0 *and an increasing function* $g(t)$ *with* $g(0) = 0$ *and* $g(t) = O(t\log(1/t)^{d+1})$ $(t \searrow 0)$ *such that if* $z_0 \in reg(\epsilon_0)$, *that is for some positive number* R_0 *and positive integer* λ_0 *possibly depending on* z_0,

$$\frac{1}{R_0^d} \int_{P_{g(R_0)(z_0)}\cap(0,T)\times\overline{\Omega}} \mathbf{e}_\lambda(z)\, dz < \epsilon_0 \qquad (3.6)$$

implies

$$\sup_{z\in P_{R_0}(z_0)\cap(0,T)\times\overline{\Omega}} \mathbf{e}_\lambda(z) \le \frac{C}{R_0^2} \qquad (3.7)$$

as long as any λ *is more than or equal to* λ_0.

Definition 3.8 *In the sequel we respectively mean* **sing** *and* **reg** *by* $\mathbf{sing}(\epsilon_0)$ *and* $\mathbf{reg}(\epsilon_0)$.

Lemma 3.9 *Pick up any point* $z_0 \in \mathbf{reg}$ *and fix it. On the parabolic cylinder* $P_{R_0/2}(z_0)$ *which is the half size of the cylinder in* (3.7)*, we obtain*

$$|u_\lambda(t,x) \,-\, u_\lambda(s,x)| \;\leq\; \frac{C}{R_0}|t-s|^{1/2} \tag{3.8}$$

for any points t *and* s *in* $[t_0 - (R_0/2)^2, t_0 + (R_0/2)^2]$ *and* $x \in \overline{B}_{R_0/2}(x_0)$ *with* $z_0 = (t_0, x_0)$*.*

Theorem 3.10 (Singular Set). *The set of* **sing** *is a closed set and*

$$\mathcal{H}^{(d)}(\mathbf{sing}) \;=\; 0 \tag{3.9}$$

holds with respect to the parabolic metric.

We finally exhibit the strong convergence of $\{u_{\lambda(\nu)}\}$ $(\nu = 1, 2, \ldots)$ to a WHHF u in $H^{1,2}$-topology as $\lambda(\nu) \nearrow \infty$. The estimates of $\mathcal{H}^{(d)}(\mathbf{sing}) = 0$ in Theorem 3.10 plays a crucial role in the proof. However since the proof is as same as in the one in K.Horihata [9, Theorem 3.12], we omit it.

Theorem 3.11 (Strong Convergent of Gradients of WHHF). *For a suitable sub-sequence of* $\{\lambda(\nu)\}$ *still denoted by* $\{\lambda(\nu)\}$ $(\nu = 1, 2, \ldots)$*, a sequence of the gradients of the GLHF,* $\{\nabla u_{\lambda(\nu)}\}$ $(\nu = 1, 2, \ldots)$ *converges strongly to the ones of the WHHF in* $L^2((0,T); L^2_B(\Omega; \mathbb{R}^{d(D+1)}))$*.*

4 Proof of Main Theorems

By utilizing a few ingredients and properties on the WHHF and the GLHF, this chapter proves Theorem 1.1, Theorem 1.2 and Theorem 1.3 in Chapter 1.

4.1 Proof of Theorem 1.1

We discuss a partial boundary regularity on the WHHF constructed above: On account of Theorem 3.10 and Theorem 3.11, we obtain

$$\mathbf{sing} \;=\; \bigcap_{R>0} \Big\{ z_0 \in (0,T) \times \overline{\Omega}\,;\; \frac{1}{2R^d} \int_{P_R(z_0) \cap Q(T)} |\nabla u|^2 \, dz \;\geq\; \epsilon_0 \Big\}, \tag{3.10}$$

$$\mathbf{reg} \;=\; (0,T) \times \overline{\Omega} \setminus \mathbf{sing}, \tag{3.11}$$

where a number ϵ_0 is a positive constant appeared in Theorem 3.7. From Theorem 3.10, we see that **sing** is closed. The size of **sing** is analogously measured as in [9, Theorem 3.10]; So does the smoothness of our WHHF on **reg**

We readily see that the WHHF u satisfies a monotonical inequality (i) and a reverse Poincaré inequality (ii) from Theorem 2.1 with Theorem 3.11 and Corollary 3.5. $\square$

4.2 Proof of Theorem 1.2

The proof is done by a direct combination of Theorem 2.1 and Theorem 3.1. $\square$

4.3 Proof of Theorem 1.3

If we recall Theorem 3.7, we have only to show that for any point $z_0 = (t_0, x_0)$,

$$\int\limits_{P_r(z_0)\cap Q(T)} |\nabla u|^2 \, dz \; < \; \frac{\epsilon_0^2 r^d}{2} \tag{3.12}$$

holds whereupon r is a positive number. If we use a stereo-graphic projection given by the mapping $v = (v_i)$ $(i = 1, 2, \ldots, D)$ of

$$\begin{cases} u^i & = \dfrac{2v^i}{1+|v|^2} \quad (i \; = \; 1, 2, \ldots, D), \\ u^D & = \dfrac{1-|v|^2}{1+|v|^2} \end{cases} \tag{3.13}$$

with $|v|^2 = \sum\limits_{i=1}^{D} (v^i)^2$, our equation (1.8) becomes for v^i $(i \; = \; 1, 2, \ldots, D)$

$$\frac{\partial}{\partial t}\Big(\frac{v^i}{1+|v|^2}\Big) \; = \; \triangle \frac{v^i}{1+|v|^2} \; + \; 4\Big|\frac{\nabla v}{1+|v|^2}\Big|^2 \frac{v^i}{1+|v|^2} \quad \text{in } Q(T). \tag{3.14}$$

By multiplying (3.14) by v^i, we have

$$\frac{\partial W(|v|^2)}{\partial t} \; - \; \triangle W(|v|^2) \; + \; \frac{4|\nabla v|^2}{(1+|v|^2)^2} \; = \; 0, \tag{3.15}$$

where the function W is given by

$$W(x) \; = \; \int_0^x \frac{(1-t^2)\, dt}{(1+t^2)^2} \quad (x \; \in \; \mathbb{R}).$$

Due to a maximal principle, we verify

$$W(|v|^2) \ \leq \ W(1-\theta_0),$$

with $\theta_0 = \mathrm{dist}(v_0(\overline{\Omega}), \partial B_1^D)/2$.

While we readily see that there exists a positive integer k_0 possibly depending on d and θ_0 such that

$$\mathrm{esssup}_{P_{d_0/2^{k_0-1}}(z_0)} W(|v|^2) \ - \ \mathrm{esssup}_{P_{d_0/2^{k_0}}(z_0)} W(|v|^2) \ < \ \epsilon_0^2$$

because of

$$\begin{aligned} &\sum_{k=1}^{\infty}(\mathrm{esssup}_{P_{d_0/2^{k-1}}(z_0)} W(|v|^2) \ - \ \mathrm{esssup}_{P_{d_0/2^{k}}(z_0)} W(|v|^2)) \\ &= \ \mathrm{esssup}_{P_{d_0}(z_0)} W(|v|^2) \ \leq \ C. \end{aligned}$$

Consequently, by employing (3.15) and a sub-linear estimate on W, we deduce

$$\int_{P_{d_0/2^k}(z_0)\cap Q(T)} |\nabla u|^2 \, dz \ < \ C\epsilon_0^2 (d_0/2^k)^d \ \leq \ \epsilon_0 (d_0/2^k)^d \tag{3.16}$$

as long as a positive number ϵ_0 is less than $1/C$.

Moreover from (3.16), an application on a ϵ-regularity theory in which we exploit (3.2) or (3.2) to our WHHF u verifies

$$\int_{P_r(z_0)\cap Q(T)} |\nabla u|^2 \, dz \ < \ Cr^{d+\alpha_0} \tag{3.17}$$

for all positive number r less than $d_0/2^{k_0+1}$, where α_0 is a positive number less than 1 and is independent of a point z_0 and a positive number r. Thus we arrive at

$$\int_{P_{g(r)}(z_0)\cap Q(T)} |\nabla u|^2 \, dz \ < \ \epsilon_0 r^d,$$

where the function g was defined in Theorem 3.7.

Here we selected a positive number r so as to be less than $(C\epsilon_0)^{1/\alpha_0}$, which conclude our claim. □

References

[1] Chen, Y.: The weak solutions to the evolution problems of harmonic maps, Math. Z. **2**01 (1989), no. 1, 69–74.

[2] Chen, Y.: Dirichlet problems for heat flow of harmonic maps in higher dimensions, Math. Z. **2**08 (1991), 557–565.

[3] Chen, Y., Lin, F.H.: Evolution of harmonic maps with Dirichlet boundary condition, Comm. Anal. Geom. **1** (1993), no. 3-4, 327–346

[4] Chen, Y., Li, J., Lin, F.H.: Partial Regularity for Weak Heat Flows into Spheres, Comm. Pure. Appl. Math, **4**8 (1995), no.4, 429–448.

[5] Cohl, S.: Fundamental Solution of Laplace's Equation in Hyperspherical Geometry, Symmetry, Integrability, Geometry and Integrability, **7** (2011), no.108 1–14.

[6] Giaquinta, M.: Multiple integrals in the calculus of variations and nonlinear elliptic problems, Annals of Mathematics Studies, **1**05 (1983), Princeton Univ. Press.

[7] Giaquinta, M., Struwe, M.: On the Partial Regularity of Weak Solutions of Nonlinear Parabolic Systems, Math. Z. **1**79 (1982), no. 4, 437–451.

[8] Hildebrandt, S.,Widman.K.-O .: Some regularity results for quasi-linear elliptic systems of second order, Math. Z. **1**42 (1975), 67-86.

[9] Horihata, K.: On a new harmonic heat flow with the reverse Hölder inequalities, Preprint.

[10] John, F.: Partial Differential Equations, Fourth edition, (1982), Springer Verlag.

[11] Jost, J., M. Meier.: Boundary regularity of minima for certain functionals, Math. Ann. **2**62, (1983), 549–561.

[12] Ladyžhenskaya, O. A., Solonnikov, V. A., Ural'ceva, N. N.: Linear and quasi-linear equations of parabolic type, Translations of Mathematical Monographs. **2**3 (1968), American Mathematical Society, Providence R.I.

[13] Poon, C.: Some new harmonic maps from B^3 to S^2 J. Diff. Geom. **4**3 (1991), 165–168.

[14] Qing, J.: Boundary Regularity of weakly harmonic maps from surfaces. J. Funct. Anal. **1**14 (1993), 458–466.

[15] Simon, L.: Theorems on Regularity and Singularity of Energy Minimizing Maps, Lectures in Mathematics ETH Zürich, Birkhäuser (1996).

[16] Schoen, R. S., Uhlenbeck, K.: Boundary regularity and the Dirichlet problem for harmonic maps, J. Differential Geom. **1**8 (1983), no. 2, 253–268.

[17] Wang, C. Y.: Boundary Partial Regularity for a class of Harmonic maps, Comm. in partial differential equations, **24** (1999), no.1, 355–368.

address 6-3, Aramaki, Aoba-ku, Sendai, Miyagi, JAPAN
e-mail horihata@m.tohoku.ac.jp